




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/53, 15/82, 15/11, 9/02, 5/10, A01H 5/00, 5/10, A23L 1/00		A1	(11) International Publication Number: WO 00/53771
			(43) International Publication Date: 14 September 2000 (14.09.00)
(21) International Application Number: PCT/US00/05915		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 8 March 2000 (08.03.00)			
(30) Priority Data: 60/123,267 8 March 1999 (08.03.99) US			
(71) Applicant (for all designated States except US): THE SAMUEL ROBERTS NOBLE FOUNDATION, INC. [US/US]; 2510 Sam Noble Parkway, Ardmore, OK 73401 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): STEELE, Christopher, L. [US/US]; 617 Campbell, Ardmore, OK 73401 (US). DIXON, Richard, A. [GB/US]; 206 Woods Lane, Ardmore, OK 73401 (US).		Published / With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(74) Agents: HANSEN, Eugenia, S. et al.; Sidley & Austin, Suite 3400, 717 N. Harwood, Dallas, TX 75201-6570 (US).			

(54) Title: GENETIC MANIPULATION OF ISOFLAVONOIDS



(57) Abstract

Soybean and *Medicago truncatula* CYP93C genes have been isolated which encode a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone. Plants can now be genetically engineered to produce isoflavones that provide potential human health benefits and increase disease resistance in plants. Isoflavones can now be produced in transgenic plants species in which isoflavones do not naturally occur, i.e., in species other than legumes. Alternatively, introducing infection-inducible isoflavonoid biosynthesis into non-legumes qualitatively complements these plants' phytoalexin defenses against microbial pathogens, whereas over-expression of the isoflavonoid pathway in legumes quantitatively increases this defense response. Finally, modifying the extend of production of isoflavonoids in legume roots positively impacts nodulation efficiency and therefore plant yield.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

GENETIC MANIPULATION OF ISOFLAVONOIDS

TECHNICAL FIELD OF THE INVENTION

The invention relates to gene manipulation in plants.

BACKGROUND OF THE INVENTION

5 The flavonoids are a major class of phenylpropanoid-derived plant natural products. Their fifteen carbon (C₆-C₃-C₆) backbone can be arranged as a 1,3-diphenylpropane skeleton (flavonoid nucleus) or as a 1,2-diphenylpropane skeleton (isoflavonoid nucleus). Although 1,3-diphenylpropane flavonoid derivatives are almost ubiquitous among terrestrial plants, the 1,2-diphenylpropane isoflavonoids are
10 restricted primarily to the Leguminosae, although they occur rarely in other families such as the Apocynaceae, Pinaceae, Compositae, and Moraceae (Tahara, S. and R. K. Ibrahim, 1995, "Prenylated isoflavonoids - an update," *Phytochemistry* 38: 1073-1094).

15 The limited taxonomic distribution of the isoflavonoids is directly related to the occurrence of the enzyme complex isoflavone synthase (IFS), which catalyzes the aryl migration reaction leading to the formation of an isoflavone from a flavanone. While flavanones are ubiquitous in higher plants, the IFS reaction, which is a two-step process specific for isoflavonoid biosynthesis (Kochs, G. and H. Grisebach, 1986, "Enzymic synthesis of isoflavones," *European J Biochem* 155: 311-318), is limited to
20 the Leguminosae and the other diverse taxa in which isoflavonoids are occasionally found.

25 The presence of isoflavonoids provides several advantages to plants. One such advantage is provided by the function of isoflavonoids as antimicrobial phytoalexins in plant-microbe interactions. For example, the simple isoflavones daidzein and genistein act as initial precursors in the biosynthesis of various antimicrobial isoflavonoid phytoalexins in a wide variety of legumes (Dixon, R. A. and N. L. Paiva, 1995, "Stress-induced phenylpropanoid metabolism," *Plant Cell* 7: 1085-1097). Isoflavonoid compounds have been shown to accumulate in infected plant cells to

levels known to be antimicrobial *in vitro*. The temporal, spatial and quantitative aspects of accumulation are consistent with a role for these compounds in disease resistance (Rahe, J. E., 1973, "Occurrence and levels of the phytoalexin phaseollin in relation to delimitation at sites of infection of *Phaseolus vulgaris* by *Colletotrichum lindemuthianum*," *Canadian J Botany* 51: 2423-2430; Hadwiger, L. A. and D. M. Webster, 1984, "Phytoalexin production in five cultivars of pea differentially resistant to three races of *Pseudomonas syringae* pv. *pisi*," *Phytopathology* 74: 1312-1314; Long, et al., 1985, "Further studies on the relationship between glyceollin accumulation and the resistance of soybean leaves to *Pseudomonas syringae* pv. *glycinea*," *Phytopathology* 75: 235-239; Bhattacharyya, M. K. and E. W. B. Ward, 1987, "Biosynthesis and metabolism of glyceollin I in soybean hypocotyls following wounding or inoculation with *Phytophthora megasperma* f. sp. *glycinea*," *Physiol and Mol Plant Pathology* 31: 387-405). Moreover, it has been reported that many plant pathogens are much more sensitive to phytoalexins of non-host species than they are to the phytoalexins of their natural hosts, because they can often detoxify the host's phytoalexins. (VanEtten, et al., 1989, "Phytoalexin detoxification: importance for pathogenicity and practical implications," *An Rev Phytopathology* 27: 143-164).

Isoflavonoids also function in plant-microbe interactions in the establishment of bacterial or fungal symbioses with plants. Isoflavonoids have been reported to regulate bacterial nodulation genes, acting as a major *nod* gene inducer (Kosslak, et al., 1987, "Induction of *Bradyrhizobium japonicum* common *nod* genes by isoflavones isolated from *Glycine max*," *Proc Natl Acad Sci USA* 84: 7428-7432) and/or transcription activator (Dakora, et al., 1993, "Common bean root exudates contain elevated levels of daidzein and coumestrol in response to *Rhizobium* inoculation," *Mol Plant-Microbe Interact* 6: 665-668). Isoflavonoids have also been shown to have a role on the establishment of the symbiotic vesicular arbuscular mycorrhizal (VAM) association of the fungus *Glomus* with legume roots. (Kape, et al., 1992, "Legume root metabolites and VA-mycorrhiza development," *J Plant Physiol* 141: 54-60). Xie et al have reported that the isoflavonoids coumestrol, daidzein and genistein have small but significant stimulatory effects on the degree of mycorrhizal colonization of soybean, and that one effect of isoflavonoids on the

soybean mycorrhizal symbiosis could be via induction of nodulation factors from co-colonizing Rhizobia, since nod-factors have also been shown to stimulate fungal colonization (Xie, et al., 1995, "Rhizobial nodulation factors stimulate mycorrhizal colonization of nodulating and nonnodulating soybeans," *Plant Physiology* 108: 1519-1525).

In addition to the advantages that the presence of isoflavonoids confers to plants, a significant body of evidence indicates that dietary consumption of isoflavonoids can provide benefits to human health. Dietary isoflavones have been ascribed strong cancer chemopreventative activity in humans, and display a range of pharmacological activities suggestive of various other health promoting effects, including phytoestrogen activity as both estrogenic and anti-estrogenic agents (Coward, et al., 1993, "Genistein, daidzein, and their -glycoside conjugates: antitumor isoflavones in soybean foods from American and Asian diets," *J Agricultural and Food Chemistry* 41: 1961-1967; Martin, et al., 1996, "Interactions between phytoestrogens and human sex steroid binding protein," *Life Sciences* 58: 429-436); anticancer effects associated with phytoestrogenic activity (Lee, et al., 1991, "Dietary effects on breast-cancer risk in Singapore," *Lancet* 337: 1197-1200; Adlercreutz, et al., 1991, "Urinary excretion of lignans and isoflavonoid phytoestrogens in Japanese men and women consuming a traditional Japanese diet," *Am J Clin Nutr* 54: 1093-1100); anticancer effects associated with inhibition of several enzymes including DNA topoisomerase and tyrosine protein kinase (Akiyama, et al., 1987, "Genistein, a specific inhibitor of tyrosine-specific protein kinases," *J Biol Chem* 262: 5592-559; Uckun, et al., 1995, "Biotherapy of B-cell precursor leukemia by targeting genistein to CD19-associated tyrosine kinases," *Science* 267: 886-891); suppression of alcohol consumption (Keung, W. M. and B. L. Vallee, 1993, "Daidzin: A potent, selective inhibitor of human mitochondrial aldehyde dehydrogenase," *Proc Natl Acad Sci USA* 90: 1247-1251; Keung, et al., 1995, "Daidzin suppresses ethanol consumption by Syrian golden hamsters without blocking acetaldehyde metabolism," *Proc Natl Acad Sci USA* 92: 8990-8993); antioxidant activity (Arora, et al., 1998, "Antioxidant activities of isoflavones and their biological metabolites in a liposomal system," *Arch Biochem Biophys* 356: 133-141; Tikkanen, et

al., 1998, "Effect of soybean phytoestrogen intake on low density lipoprotein oxidation resistance," *Proc Natl Acad Sci USA* 95: 3106-3110); effects on calcium metabolism, some of which may be linked to protective effects against osteoporosis (Tomonaga, et al., 1992, "Isoflavonoids, genistein, PSI-tectorigenin, and orobol, increase cytoplasmic free calcium in isolated rat hepatocytes," *Biochem Biophys Res Com* 182: 894-899; Draper, et al., 1997, "Phytoestrogens reduce bone loss and bone resorption in oophorectomized rats," *J Nutr* 127: 1795-1799); and cardiovascular effects (Wagner, et al., 1997, "Dietary soy protein and estrogen replacement therapy improve cardiovascular risk factors and decrease aortic cholesteryl ester content in ovariectomized cynomolgus monkeys," *Metabolism - Clinical and Experimental* 46: 698-705).

At present, the only dietary sources of isoflavonoids for humans are certain legumes such as soybean or chickpea. The development of methods to genetically manipulate isoflavonoids in plants, either to widen the source of dietary isoflavonoids for humans, or to exploit the biological activities of isoflavonoids for plant protection and improvement, is wholly dependent on the availability of cloned genes encoding the various enzymes of isoflavonoid biosynthesis. Of these, the isoflavone synthase (IFS) complex constitutes the first committed reactions, and as such represents the means to introduce isoflavonoids into plants that do not possess the pathway.

In 1984, Hagmann and Grisebach provided the first evidence for the enzymatic conversion of flavanone to isoflavone (the IFS reaction) in a cell free system (Hagmann, M. and H. Grisebach, 1984, "Enzymatic rearrangement of flavanone to isoflavone," *FEBS Letters* 175: 199-202). They demonstrated that microsomes from elicitor-treated soybean cell suspension cultures could catalyze the conversion of 2(S)-naringenin to genistein, or of 2(S)-liquiritigenin to daidzein, in the presence of NADPH. The crude microsomal enzyme preparation, which was stable at -70°C but had a half-life of only 10 minutes at room temperature, was absolutely dependent on NADPH and molecular oxygen. It was subsequently shown that the reaction proceeded in two steps. The flavanone was converted in a cytochrome P450-catalyzed reaction requiring NADPH and O₂ to the corresponding 2-hydroxyisoflavanone. This

relatively unstable compound, which could, however, be identified by mass spectrometric analysis, then underwent dehydration to yield the isoflavone. The dehydration reaction appeared to be catalyzed by an enzyme present predominantly in the cytoplasmic supernatant, although it was not possible to remove all this activity from the microsomes. The corresponding 2-hydroxyisoflavanone spontaneously converted to genistein, for example, in methanol at room temperature. Kinetic analysis indicated that the 2-hydroxyisoflavanone was formed prior to genistein, consistent with its being an intermediate in isoflavone formation. (Kochs, G. and H. Grisebach, 1986, "Enzymic synthesis of isoflavones," *European J Biochem* 155: 311-318).

Involvement of cytochrome P450 in the 2-hydroxyisoflavanone synthase reaction was confirmed by inhibition by CO, replacing O₂ with N₂, and examining the effects of a range of known P450 inhibitors of which ancymidol was the most effective. The enzyme co-migrated with the endoplasmic reticulum markers cinnamate 4-hydroxylase (another cytochrome P450) and cytochrome b5 reductase on Percoll gradients. The enzyme is stereoselective, and (2*R*)-naringenin is not a substrate. (Kochs, G. and H. Grisebach, 1986, "Enzymic synthesis of isoflavones," *European J Biochem* 155: 311-318).

The origin of the 2-hydroxyl group was determined from studies on the IFS present in microsomes from elicited cell cultures of *Pueraria lobata*. ¹⁸O from ¹⁸O₂ was incorporated into the 2-hydroxyl group, resulting in a 2-hydroxyisoflavanone with molecular ion shifted by two mass units, whereas there was no corresponding shift in the molecular ion of daidzein, consistent with the subsequent dehydration reaction (Hashim, et al., 1990, "Reaction mechanism of oxidative rearrangement of flavanone in isoflavone biosynthesis," *FEBS Letters* 271: 219-222). The currently accepted model for the reaction pathway of IFS as illustrated in Fig. 1, therefore, involves P450-catalyzed hydroxylation coupled to aryl migration, a reaction with mechanistic similarities to the well described proton migration mechanism of some P450 reactions (Hakamatsuka, et al., 1991, "P-450-dependent oxidative rearrangement in isoflavone

biosynthesis: reconstitution of P-450 and NADPH:P450 reductase," *Tetrahedron* 47: 5969-5978).

Currently, there have been no reports on purification to homogeneity or molecular cloning of the cytochrome P450 of the IFS complex because of the extreme lability of the enzyme. The 2-hydroxyisoflavanone synthase cytochrome P450 from *Pueraria* has been solubilized with Triton X-100, and partially purified by DEAE-Sephacrose chromatography; the enzymatic reaction could be reconstituted by addition of NADPH cytochrome P450 reductase that separated from the hydroxylase on the ion exchange column (Hakamatsuka, et al., 1991, *Tetrahedron* 47: 5969-5978). A 2-hydroxyisoflavanone dehydratase has been purified from elicitor-treated *P. lobata* cells, and has been shown to be a soluble monomeric enzyme of subunit Mr 38,000 (Hakamatsuka, et al., 1998, "Purification of 2-hydroxyisoflavanone dehydratase from the cell cultures of *Pueraria lobata*," *Phytochemistry* 49: 497-505). It is not yet clear whether this enzyme physically associates with the P450 hydroxylase catalyzing the aryl migration, or even whether this activity is essential for isoflavone formation *in planta* in view of the spontaneous conversion of 2-hydroxyisoflavanone to isoflavone.

Flavanone is a potential substrate for more than one type of hydroxylation reaction at the 2-position. Thus, elicitor-treated cell cultures of alfalfa and *Glycyrrhiza echinata* have been shown to accumulate the dibenzoylmethane licodione (Kirikae, et al., 1993, "Biosynthesis of a dibenzoylmethane, licodione, in cultured alfalfa cells induced by yeast extract," *Biosci Biotech Biochem* 57: 1353-1354). Licodione synthase is, by classical criteria, a cytochrome P450, the activity of which is induced by yeast elicitor in *Glycyrrhiza* cells (Otani, et al., 1994, "Licodione synthase, a cytochrome P450 monooxygenase catalyzing 2-hydroxylation of 5-deoxyflavanone, in cultured *Glycyrrhiza echinata* L. cells," *Plant Physiol* 105: 1427-1432). The reaction it catalyzes involves 2-hydroxylation of flavanone followed by hemiacetal opening instead of aryl migration, and the reaction was thought to have mechanistic similarities to the flavone synthase II enzyme previously characterized from soybean (Kochs, G. and H. Grisebach, 1987, "Induction and characterization of a NADPH-dependent flavone synthase from cell cultures of soybean," *Z. Naturforsch* 42C: 343-

348). A gene encoding the flavone synthase II/licodione synthase from *Glycyrrhiza* has been cloned (Akashi, et al., 1998, "Identification of a cytochrome P450 cDNA encoding (2S)-flavanone 2-hydroxylase of licorice (*Glycyrrhiza echinata* L.: *Fabaceae*) which represents licodione synthase and flavone synthase II," *FEBS Letters* 5 431: 287-290), and a different cytochrome P450 gene encoding flavone synthase II has recently been cloned from *Gerbera hybrida* (Martens, S. and G. Forkmann, "Cloning and expression of flavone synthase II from *Gerbera* hybrids," *Plant J* 20: 611-618).

Although the reactions catalyzed by IFS are critical for the formation of all isoflavonoids in plants, there have been no previous reports of the isolation of genes encoding components of isoflavone synthase, although genes encoding most of the 10 other enzymes of the isoflavonoid pathway, including downstream enzymes converting simple isoflavones to antimicrobial phytoalexins, have been characterized (Dixon, et al., 1995, "The isoflavonoid phytoalexin pathway: from enzymes to genes to transcription factors," *Physiologia Plantarum* 93: 385-392). Thus, the 15 unavailability of isoflavone synthase genes has made it heretofore impossible to utilize the downstream genes for regulating isoflavonoid concentrations in legumes and other plants that do have the isoflavonoid pathway, or for engineering antimicrobial and pharmacologically active isoflavonoids in transgenic plants of species that do not have the isoflavonoid pathway.

20 Genes encoding the enzyme catalyzing the first step of the isoflavone synthase reaction have now been isolated and purified from soybean and *Medicago truncatula* (barrel medic).

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 depicts the currently accepted model for the reaction pathway of IFS 25 wherein the flavanone is converted in a cytochrome P450-catalyzed reaction requiring NADPH and O₂ to the corresponding 2-hydroxyisoflavanone which then undergoes dehydration to yield the isoflavone.

Fig. 2 depicts the nucleotide sequence of soybean CYP93C1v2.

Fig. 3 depicts the amino acid sequence of soybean CYP93C1v2 compared to licorice CYP93B1.

Fig. 4 depicts the nucleotide sequence of *Medicago truncatula* mtIFSE3.

Fig. 5 depicts the amino acid sequence of *Medicago truncatula* mtIFSE3
5 compared to soybean CYP93C1v2.

Fig. 6A and Fig. 6B depict HPLC traces of extracts from pooled tissues (leaves, shoots, flowers) of *Arabidopsis thaliana* ecotype Columbia harboring an empty tDNA vector (Fig. 6A) and *Arabidopsis thaliana* ecotype Columbia harboring the soybean CYP93C1v2 cDNA sequence (Fig. 6B). The empty vector transformed
10 line contains a number of flavonol glycosides and other phenolic compounds that are also present in the CYP93C1v2 transformed line. These compounds were identified as (a) rhamnose (Rha)- glucose (Glc)- quercetin (Q), (b) uncharacterized conjugate of Q, (c) Rha-Glc-Rha-Kaempferol (K), (d) Glu-Rha-Q, (e) Rha-Rha-Q, (f) Glc-Rha-K, (g) sinapic acid, (h) Rha-Rha-K. Three additional compounds were observed in the
15 CYP93C1v2 transformed line (Fig. 6B), and labeled "1," "2" and "3." Fig. 6C depicts a total ion chromatogram of partially purified peaks 2 and 3, and the insets show the specific ions generated from these compounds. Peak 2 has a parental molecular mass ion of 579.5 consistent with genistein conjugated to a glucose-rhamnose disaccharide, and two further mass ions of 417.5 and 271.3, representing Rha-genistein and free
20 genistein, respectively. Peak 3, which has a parental molecular ion of mass 417.5, is thereby identified as Rha-genistein.

Fig. 7A and Fig. 7B depict HPLC traces of the same extracts as shown in Fig. 6A (empty-vector transformed) and Fig. 6B, (CYP93C1v2 transformed), but following digestion with β -glucosidase. Peaks 2 and 3 remained at the same retention time as in
25 Fig. 6A and 6B. However, Peak 1 disappeared, and was replaced with a new Peak 4 of much later retention time. Fig. 7C shows the total ion chromatograph of purified Peak 4, and the inset shows the parental molecular ion, with mass of 271.2, consistent with Peak 4 being free genistein. Fig. 7D shows a total ion chromatograph, and the parental molecular ion, of an authentic sample of genistein.

Fig. 8A , 8B, 8C and 8D are high performance liquid chromatography (HPLC) chromatograms depicting the presence of new peaks at RT 29.96 and 37.7 min representing the presence of the isoflavone daidzein formed from the flavanone liquiritigenin, or the isoflavone genistein formed from the flavanone naringenin, in insect cell microsomes expressing CYP93C1v2. Fig. 8A depicts the presence of NADPH during incubation with liquiritigenin. Fig. 8B depicts the absence of NADPH during incubation with liquiritigenin. Fig. 8C depicts the presence of NADPH during incubation with naringenin. Fig. 8D depicts the lack of a reaction when soybean CYP93E expressed in insect cells is incubated with liquiritigenin in the presence of NADPH.

Fig. 9A and Fig. 9B are mass spectra of BSTFA (*N,O*-bis(trimethylsilyl) trifluoroacetamide) derivatives. Fig. 9A depicts the mass spectrum of the BSTFA derivative of the product of the reaction catalyzed by CYP93C1v2 in insect cells using liquiritigenin as substrate, and Fig. 9B shows the mass spectrum of the BSTFA derivative of an authentic sample of daidzein .

SUMMARY OF THE INVENTION

In one aspect, the invention is a method for introducing into a naturally non-isoflavonoid-producing plant species the enzyme catalyzing the aryl migration of a flavanone to form an isoflavanone intermediate or an isoflavone, comprising introducing a DNA segment encoding the enzyme into the plant to form a transgenic plant, wherein the transgenic plant expresses the DNA segment under the control of a suitable constitutive or inducible promoter when the transgenic plant is exposed to conditions which permit expression. The DNA segment can comprise isolated genomic DNA or recombinant DNA. Preferably, the DNA segment is a CYP93C gene. An exemplary DNA segment from a soybean CYP93C gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another preferred DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene, more preferably, the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4. Plants transformed by this method may also preferably express chalcone synthase,

chalcone reductase, and chalcone isomerase genes to cause in vivo formation of daidzein or a daidzein derivative, and the chalcone synthase, chalcone reductase, and chalcone isomerase genes may also be transgenes. Plants transformed by this method may also preferably further comprise downstream genes, for example, isoflavone *O*-methyltransferase, isoflavone 2'-hydroxylase, isoflavone reductase, and vestitone reductase, to metabolize a formed isoflavone to biologically active isoflavonoid derivatives or conjugates. The plant can comprise isoflavone 4'-*O*-methyl-transferase to cause formation of biochanin A or a biochanin A derivative from the isoflavanone intermediate. An exemplary flavanone substrate for this transformation method is liquiritigenin and/or naringenin.

In another aspect, the present invention is a method for increasing the level of isoflavonoid compounds in naturally isoflavonoid-producing plants comprising introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to yield an isoflavonoid to form a transgenic plant, wherein the transgenic plant expresses the DNA segment under the control of a suitable constitutive or inducible promoter when the transgenic plant is exposed to conditions which permit expression. With this method, the resulting isoflavonoid can be an isoflavanone intermediate, an isoflavone, an isoflavone derivative, and an isoflavone conjugate. The DNA segment can comprise isolated genomic DNA or recombinant DNA. Preferably, the DNA segment is a CYP93C gene. An exemplary DNA segment from a soybean CYP93C gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another preferred DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene, more preferably, the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4. An exemplary flavanone substrate for this transformation method is liquiritigenin and/or naringenin.

In another aspect, the invention is a method for synthesizing an isoflavanone intermediate or an isoflavone from a flavanone by expressing a recombinant CYP93C gene segment in a suitable bacterial, fungal, algal, or insect cell system. An exemplary gene segment consists essentially of the sequence from about nucleotide 36

to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another exemplary gene segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

In another aspect, the invention is a method of reducing the levels of isoflavonoid compounds in a naturally isoflavonoid-producing plant comprising introducing and expressing an antisense or gene silencing construct that contains an intact CYP93C gene or segments thereof into the plant. An exemplary gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another exemplary gene consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

In another aspect, the invention is a naturally non-isoflavonoid-producing plant cell transformed by introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to form an isoflavanone intermediate or an isoflavone, wherein the transformed plant cell expresses the DNA segment under the control of a suitable constitutive or inducible promoter when exposed to conditions which permit expression. The DNA segment can comprise isolated genomic DNA or recombinant DNA. Preferably, the DNA segment is a CYP93C gene. An exemplary DNA segment from a soybean CYP93C gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another preferred DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene, more preferably, the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4. Plants transformed by this method may also preferably express chalcone synthase, chalcone reductase, and chalcone isomerase genes to cause in vivo formation of daidzein or a daidzein derivative, and the chalcone synthase, chalcone reductase, and chalcone isomerase genes may also be transgenes. Plants transformed by this method may also preferably further comprise downstream genes, for example, isoflavone *O*-methyltransferase, isoflavone 2'-hydroxylase, isoflavone reductase, and vestitone reductase, to metabolize a formed isoflavanone intermediate to biologically active isoflavonoid

derivatives or conjugates. The plant can comprise isoflavone 4'-O-methyl-transferase to cause formation of biochanin A or a biochanin A derivative from the isoflavanone intermediate.

5 In another aspect, the invention is a naturally isoflavonoid-producing plant cell transformed by introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to yield an isoflavonoid to form a transformed plant cell, wherein the transformed plant cell expresses the DNA segment under the control of a suitable constitutive or inducible promoter when exposed to conditions which permit expression. With this method, the resulting isoflavonoid can be an isoflavanone
10 intermediate, an isoflavone, an isoflavone derivative, and an isoflavone conjugate. The DNA segment can comprise isolated genomic DNA or recombinant DNA. Preferably, the DNA segment is a CYP93C gene. An exemplary DNA segment from a soybean CYP93C gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another
15 preferred DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene, more preferably, the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

In another aspect, the invention is a transgenic plant cell having reduced levels of isoflavonoid compounds, the plant cell transformed by introducing an antisense or
20 gene silencing construct that contains an intact CYP93C gene or segments thereof into the plant cell. An exemplary gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. Another exemplary gene consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

25 In another aspect, the invention is an isolated gene or DNA segment comprising a portion which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the portion consists essentially of about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1. An exemplary gene is the soybean
30 gene encoding the enzyme catalyzing the aryl migration of liquiritigenin. Another

exemplary gene is the soybean gene encoding the enzyme catalyzing the aryl migration of naringenin.

5 In another aspect, the invention is a protein encoded by a portion of an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the portion consists essentially of about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

10 In another aspect, the invention is an isolated gene or DNA segment comprising a portion which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the portion is a *Medicago truncatula* homolog of a CYP93C gene. An exemplary gene or DNA segment consists essentially of about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4. An exemplary gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of
15 liquiritigenin. Another exemplary gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of naringenin.

In another aspect, the invention is a protein encoded by a portion of an isolated gene or a DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone,
20 wherein the portion is a *Medicago truncatula* homolog of a CYP93C gene.

In yet another aspect, the invention is a food comprising edible transgenic plant material capable of being ingested for its nutritional value, wherein the transgenic plant has been transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an
25 isoflavanone intermediate or an isoflavone, and wherein the transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a food comprising at least one isoflavonoid, wherein the isoflavonoid is isolated from a transgenic plant transformed

with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, and wherein the transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a composition comprising at least a portion of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment, and wherein the composition is suitable for ingestion as a food stuff, a nutritional supplement, an animal feed supplement, or a nutraceutical.

In yet another aspect, the invention is a composition comprising an isoflavonoid suitable for administration as a food stuff, a nutritional supplement, an animal feed supplement, a nutraceutical, or a pharmaceutical, wherein the isoflavonoid is isolated from at least a portion of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, and wherein the transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a method of increasing the nutritional value of a plant by transforming the plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a method of using a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment, to provide a nutraceutical benefit to a human or animal administered the isoflavonoid. The isoflavonoid can be administered by ingestion of at least a portion of the plant. The isoflavonoid can also be administered by ingestion of a composition comprising an isoflavonoid isolated from the plant.

In yet another aspect, the invention is a method of using an isoflavonoid isolated from a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment, to provide a pharmaceutical benefit to a patient administered the isoflavonoid.

In yet another aspect, the invention is a method of increasing disease resistance in a plant by transforming the plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a method of increasing nodulation efficiency of a leguminous plant by transforming the plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid released from the roots

when compared to the level of the isoflavonoid released from the roots of plants of the same species which do not comprise the isolated gene or DNA segment.

5 In yet another aspect, the invention is a transgenic leguminous plant exhibiting increased nodulation efficiency transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid released from the roots when compared to the level of the isoflavonoid released from the roots of plants of the same species which do not comprise the isolated gene or DNA segment.

10 In yet another aspect, the invention is a method of increasing bacterial or fungal symbiosis in a plant by transforming the plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the
15 level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a transgenic plant exhibiting increased bacterial or fungal symbiosis transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone
20 to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

In yet another aspect, the invention is a transgenic plant comprising at least one
25 recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the recombinant DNA sequence.

In yet another aspect, the invention is seed from a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid
5 when compared to the level of the isoflavonoid in plants of the same species which do not comprise the recombinant DNA sequence.

In yet another aspect, the invention is progeny from a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or
10 an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the recombinant DNA sequence.

In yet another aspect, the invention is progeny from seed of a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that
15 can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the recombinant DNA sequence.

In yet another aspect, the invention is use of a transgenic plant transformed
20 with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment, for the preparation of a nutraceutical
25 preparation for achieving a nutritional effect.

In yet another aspect, the invention is use of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein the transgenic plant exhibits an increased level of an isoflavonoid

when compared to the level of the isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment, for the preparation of a pharmaceutical preparation for achieving a therapeutic effect.

DETAILED DESCRIPTION

5 One aspect of the present invention is an isolated gene which encodes the first step of the isoflavone synthase reaction: a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavone. Genes and corresponding cDNA of the soybean or *Medicago truncatula* CYP93 family have been isolated. The enzymes encoded by the genes of the present invention are isoflavone synthases (IFS) and can
10 catalyze the aryl migration of a flavanone to yield an isoflavone either directly or through the intermediacy of a 2-hydroxyisoflavanone. One isolated soybean gene is classified as *CYP93C1v2*.

 Cytochrome P450 enzymes belong to a large superfamily of enzymes that are abundant in every living organism. The P450 nomenclature committee has
15 determined that each P450 should carry a "CYP" designation and arbitrarily divided the superfamily into families (alphabetical designation), subfamilies (numerical designation) and allelic variants ("v" plus numerical designation) based on amino acid identity of >40%, >55%, and >97%, respectively (Nelson, et al. 1993. "The P450
20 superfamily update on new sequences, gene mapping, accession numbers, early trivial names of enzymes, and nomenclature," *DNA Cell Biol* 12:1). Thus, CYP93C1v2 is a variant of the first described P450 belonging to the third subfamily (C) of the ninety-third P450 family.

 Utilizing the procedures presented herein, any plant known to produce isoflavonoids may also serve as sources of suitable DNA, or coding sequences may be
25 synthesized in vitro based on the sequences for the IFS genes of the present invention. CYP93 family members can also be obtained from other plant species by polymerase chain reaction amplification methods known to those skilled in the art, using primer sequences corresponding to regions of nucleotide conservation between CYP93 family members. Furthermore, the genes of the present invention are defined by their

catalytic activity: the aryl migration of a flavanone to yield an isoflavone. The gene sequences presented as SEQ ID NO:1 and SEQ ID NO:4 are exemplary, and it is understood that modifications to these genes which do not alter the catalytic activity of its encoded protein fall within the scope of the present invention. While a preferred

5 IFS gene contains the entire open reading frame, portions of or the entire 5' and 3' untranslated regions as well as portions of the vector sequence can also be present. With the isolation and functional identification of these isoflavone synthase (IFS) genes that encode the first key step in isoflavone formation, the aryl migration reaction, it is now possible to introduce the isoflavonoid pathway into all plant species,

10 including those that do not naturally possess this pathway.

Another aspect of the present invention is a genetically modified plant which has been transformed with a gene of the present invention. For example, when the *CYP93C1v2* gene is transferred into the model plant *Arabidopsis thaliana*, which does not naturally produce isoflavonoids, the isoflavone genistein accumulates as a series of

15 glycoconjugates (Example 1). This demonstrates that the genes of the present invention can be genetically engineered into plants which do not naturally contain the isoflavonoid pathway, and the transgenic plants can then produce isoflavonoids, resulting in plants with improved disease resistance and/or value added health benefits for humans. In the present invention, unless otherwise stated, as used herein, the term

20 "plant" or "progeny" includes plant parts, plant tissue, plant cells, plant protoplasts, plant cell tissue cultures from which plants can be regenerated, plant calli, plant clumps, explants, plant cells that are intact in plants, or parts of plants, such as embryos, pollen, ovules, flowers, capsules, stems, leaves, seeds, roots, root tips, and the like. Furthermore, the present invention includes the IFS genes expressed in

25 various parts of the plant, e.g., in aerial portions of the plant useful for increasing disease resistance or production of health promoting isoflavonoid nutraceuticals, in seeds useful for increasing levels of isoflavones and their conjugates, or in roots useful for increasing disease resistance or production of nodulation gene inducing isoflavones.

In another aspect, the present invention is a method of improving disease resistance and a transgenic plant with increased disease resistance. By transforming a plant which does not naturally make isoflavones with an IFS gene of the present invention, disease resistance can be genetically engineered into the plant by providing the necessary enzyme to convert its natural flavanones into isoflavonoids. The introduction and subsequent expression of an IFS gene of the present invention into a crop species which naturally possesses the isoflavonoid pathway results in increased levels of the isoflavonoid defense compounds.

In another aspect, the present invention is a method of increasing levels of isoflavonoids that might be beneficial to the establishment of bacterial or fungal symbioses with plants and a transgenic plant with an increased capacity for symbiotic association with bacteria or fungi. Bacterial nodulation can be stimulated in transgenic leguminous plants by expression of an IFS gene of the present invention and decreased by expression of antisense constructs or constructs designed to promote gene silencing that contain an intact IFS gene or segments thereof. Mycorrhizal colonization of leguminous plants can also be increased through the introduction and expression of an IFS gene of the present invention.

In yet another aspect, the present invention is a method of producing isoflavonoid compounds in plants or any other organism to be used in nutraceuticals or pharmaceuticals to confer human or animal health benefits. Edible transgenic plants high in isoflavonoids can be utilized as food for humans and animals. Edible compositions high in isoflavonoids can also be made by incorporation of the transgenic plants or plant materials, or by incorporation of isoflavonoids isolated from the transgenic plants. Compositions useful for administration as a food stuff, a nutritional supplement, an animal feed supplement, a nutraceutical, or a pharmaceutical can be made by incorporation of the transgenic plants or plant materials, or by incorporation of isoflavonoids isolated from the transgenic plants. The nutritional value of a plant can be increased by transforming the plant with an IFS gene of the present invention and, as a result, accumulating high amounts of isoflavonoids in the plant.

The soybean IFS gene of the present invention was isolated and purified according to the detailed procedures outlined in Example 2. The DNA sequence is shown in SEQ ID NO:1 and Fig. 2, and the encoded protein sequence of the isolated soybean CYP93C clone is shown in SEQ ID NO:2 and Fig. 3. For comparison, Fig. 3 also shows the protein sequence alignment between the isolated CYP93C clone (SEQ ID NO:2) and CYP93B1 (SEQ ID NO:3), the licorice licodione synthase.

The DNA and protein sequences of the soybean CYP93C1 open reading frame were deposited in the Genbank data base under accession # AF022462. The deposition was made by Siminszky, Dewey and Corbin, and the sequence described as representing a gene induced in soybean in response to herbicide safeners. However, the function of the gene was not known and there was no understanding that it could be involved in isoflavonoid biosynthesis at the time the deposit was made (Siminszky, B., Corbin, F.T., Ward, E.R., Fleischmann, T.J. and Dewey, R.E., 1999, "Expression of a soybean cytochrome P450 monooxygenase cDNA in yeast and tobacco enhances the metabolism of phenylurea herbicides." *Proc. Natl. Acad. Sci. USA* 96: 1750-1755). The sequence of the clone characterized herein differs from CYP93C1 in three nucleotide substitutions in the open reading frame that change proline 140 to leucine, threonine 156 to isoleucine, and glutamate 295 to lysine. Thus, the soybean gene identified herein has been classified as CYP93C1v2.

The cDNA insert from CYP93C1v2 was used to probe 240,000 phage plaques from a *Medicago truncatula* root cDNA library (van Buuren, M.L., I.E. Maldonado-Mendoza, A.T. Trieu, L.A. Blaylock, and M.J. Harrison, 1999, "Novel genes induced during an arbuscular mycorrhizal (AM) symbiosis formed between *Medicago truncatula* and *Glomus versiforme*," *Mol. Plant-Microbe Interact.* 12, 171-181). Five positive plaques were purified, in vivo excised, and sequenced. A full length clone designated mtIFSE3 was completely sequenced on both strands, and shown to encode the *Medicago truncatula* homolog of soybean CYP93C1. The nucleotide sequence of mtIFSE3 is shown in SEQ ID NO:4 and Fig. 4, and the protein sequence, in SEQ ID NO:5. An alignment between the protein sequences of mtIFSE3 and CYP93C1v2 is shown in Fig. 5.

An IFS gene of the soybean or *Medicago truncatula* CYP93C subfamily or corresponding cDNA sequence, the open reading frame of which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavone, either directly or through the intermediacy of a 2-hydroxyisoflavanone, can be used to introduce the isoflavonoid pathway into any plant species that does not naturally possess this pathway. Soybean CYP93C1v2 acts on the flavanones liquiritigenin to yield daidzein, and naringenin to yield genistein. Liquiritigenin is only formed in plants that possess the enzyme chalcone reductase (CHR) (Welle, R. and Grisebach, H., 1989, "Phytoalexin synthesis in soybean cells: elicitor induction of reductase involved in biosynthesis of 6'-deoxychalcone." *Arch Biochem Biophys* 272: 97-102), and a form of chalcone isomerase that is active against 2',4,4'-trihydroxychalcone, the product of the co-action of chalcone synthase (CHS) with CHR (Dixon, R.A., Blyden, E.R., Robbins, M.P., van Tunen, A.J. and Mol, J.N.M., 1988, "Comparative biochemistry of chalcone isomerases." *Phytochemistry* 27: 2801-2808). Such genes are common in legumes, but not in most other plant families. Thus, to form daidzein in transgenic plants that do not possess the isoflavonoid pathway, it would be necessary to introduce three new genes, namely CHR, to co-act with CHS to form 2',4,4'-trihydroxychalcone, a suitable CHI to convert 2',4,4'-trihydroxychalcone to liquiritigenin, and IFS, assuming that the 2-hydroxyisoflavanone intermediate can spontaneously dehydrate in planta, a phenomenon that is demonstrated below. Without CHR present, no liquiritigenin would be formed, and IFS would only be able to act on naringenin to yield, assuming spontaneous dehydration of the 2-hydroxyisoflavanone, genistein.

The IFS genes of the present invention can be introduced into non-leguminous plants such as by standard *Agrobacterium*-based or biolistic transformation procedures (Horsch, et al., 1985, "A simple and general method for transferring genes into plants," *Science* 227:1229-1231; and Klein, et al., 1988, "Stable genetic transformation of intact *Nicotiana* cells by the particle bombardment process," *Proc Natl Acad Sci USA* 85:8502-8505). Both procedures require the construction of a plasmid vector containing a desirable transcriptional promoter driving expression of the gene of interest (in this case IFS), followed by a transcriptional terminator and a

selectable marker gene for resistance, such as to an antibiotic or a herbicide. The biolistic procedure coats metal particles with plasmid DNA containing the gene of interest and places them on a micro carrier disk. Using the biolistic apparatus, the particles are physically propelled into plant tissue. The plant tissue is then put under selection (e.g., antibiotic or herbicide) followed by regeneration. The two *Agrobacterium*-based procedures are “in planta” and “ex-planta”, respectively. Both procedures require the above gene construct to be placed into a T-DNA vector, which is then transferred into *Agrobacterium tumefaciens*. The in planta procedure places the transformed *Agrobacterium* in the presence of plant material (flower or meristem) and the plants are allowed to seed followed by selection (e.g., antibiotic or herbicide) during germination. The ex-planta procedure also places *Agrobacterium* in the presence of plant material (callus, cell culture, leaf disk, hypocotyl) which is placed directly under selection (e.g., antibiotic or herbicide) followed by regeneration.

Thus, the isoflavonoid pathway can be introduced into any plant species that does not possess the enzyme catalyzing the IFS reaction by expressing the IFS gene in transgenic plants under the control of a suitable constitutive or inducible promoter.

Example 1: Transformation of *Arabidopsis thaliana* with Soybean CYP93C1V2

Soybean CYP93C1v2 cDNA was placed in the binary plant transformation vector pCHF3, in which it is under control of the cauliflower mosaic virus 35S promoter, using standard recombinant DNA methods (Sambrook, et al. 1989. *Molecular Cloning. A Laboratory Manual*, 2nd Ed, Cold Spring Harbor Laboratory Press, New York). The gene was then transformed into plants of the crucifer, *Arabidopsis thaliana* ecotype Columbia, using *Agrobacterium tumefaciens* and a standard floral infiltration procedure (Clough, S.J. and Bent, A.F., 1998, “Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*”. *Plant J* 16: 735-743). Transgenic plants were selected by germinating the seedlings on kanamycin, and those surviving selection were allowed to set seed. T₂ seedlings expressing CYP93C1v2 were identified by standard DNA and RNA gel blot analysis (Sambrook, et al. 1989. *Molecular cloning. A Laboratory Manual*, 2nd Ed,

Cold Spring Harbor Laboratory Press, New York), and analyzed for accumulation of genistein in leaves by HPLC analysis, according to a method developed to profile the flavonoid components of *Arabidopsis* leaves (Graham, T.L., 1998, "Flavonoid and flavonol glycoside metabolism in *Arabidopsis*". *Plant Physiol Biochem* 36: 135-144).

5 Figure 6A shows a typical HPLC trace of a leaf extract from an untransformed plant. The major components are glycosides (containing glucose and rhamnose) of the flavonols kaempferol and quercetin. Plants harboring the soybean CYP93C1v2 gene showed an additional three peaks on HPLC analysis (Fig. 6B), indicated by the arrows labeled as "1," "2" and "3." No free genistein, free 2-hydroxyisoflavanone or 2-
10 hydroxyisoflavanone conjugates were observed. However, following treatment of extracts with almond β -glucosidase (Fig 7B), one of the new peaks disappeared, and free genistein was now observed, consistent with the peak being a glucoside of genistein. LC-MS analysis confirmed the identities of the new compounds as a
15 glucoside of genistein, glucose-rhamnose-genistein, and rhamnose-genistein (Figs. 6C and 7C and 7D, insets). Therefore, expression of CYP93C1v2 in transgenic *Arabidopsis* leads to formation of genistein with no requirement for an enzyme to catalyze the dehydration of the presumed 2-hydroxyisoflavanone intermediate. *Arabidopsis* plants then modify the genistein by exactly the same chemistry they use to conjugate their endogenous flavonols, namely by conjugation to glucose and
20 rhamnose. Transgenic production of conjugates of genistein are suitable for nutraceutical applications, because genistein is also glycosylated in soybean, its natural dietary source (Graham, T.L., 1991, "Flavonoid and isoflavonoid distribution in developing soybean seedling tissues and in seed and root exudates." *Plant Physiol* 95: 594-603).

25 In addition to introducing the isoflavonoid pathway into plants that do not possess this pathway, the level of isoflavonoid compounds can be controlled in plants that do possess the pathway by manipulating the level of expression of the IFS gene. Increasing the levels of isoflavonoid compounds in leguminous plants by expression of the IFS gene of the present invention in transgenic plants under the control of a
30 suitable constitutive or inducible promoter can be accomplished by standard methods

such as *Agrobacterium*-based or biolistic transformation methods known in the art. Alternatively, the level of isoflavonoid compounds in plants can be reduced by expression of antisense constructs or constructs designed to promote gene silencing that contain an intact IFS gene, or segments thereof, in transgenic plants using methods known in the art. (Bourque, J.E., 1995, "Antisense strategies for genetic manipulations in plants," *Plant Science* 105:125-149; and Angell, S. M. and D. C. Baulcombe, 1997, "Consistent gene silencing in transgenic plants expressing a replicating potato virus X RNA," *EMBO J* 16:3675-3684). Antisense constructs for gene silencing are constructed by placing the whole or part of the cDNA in a three prime to five prime orientation behind a desirable transcriptional promoter and ahead of a transcriptional terminator in a plasmid vector. The vector may be used for biolistic transformation or the new antisense gene may be transferred to a T-DNA vector for *Agrobacterium*-based transformation. The actual mechanism of silencing by antisense constructs is unknown. Homology-dependent gene silencing or co-suppression requires the over-expression of a homologous gene; therefore, to achieve co-suppression a construct is made using a strong promoter, the gene of interest (in this case IFS) and a transcriptional terminator. The gene should be transferred to plants as described above. Gene silencing is an epigenetic phenomenon that may or may not occur with a particular gene construct. When it does occur, the inhibition of gene expression can be greater than with the antisense approach.

Isoflavones can be synthesized from flavanones, utilizing recombinant IFS expressed in any suitable bacterial, fungal, algal, or insect cell system. For example, naringenin is extracted in large amounts from grapefruits. A CYP93C1 enzyme can be used convert naringenin to 2,5,7,4'-tetrahydroxyisoflavanone, which spontaneously converts to the valuable nutraceutical genistein under weak acid conditions. Furthermore, daidzin can be synthesized from liquiritigenin utilizing recombinant CYP93C1 and an isoflavone glucosyltransferase (Köster, J. and W. Barz, 1981, "UDP-Glucose: isoflavone 7-O-glucosyltransferase from roots of chick pea (*Cicer arietinum* L.)." *Arch Biochem Biophys* 212: 98-104).

Example 2: Methodology Used to Isolate and Identify IFS cDNA Clones

In an attempt to obtain cDNA clones encoding IFS, a functional genomics approach was followed. IFS activity is present in soybean seeds, which accumulate daidzein and genistein. Furthermore, IFS activity can be induced in soybean tissues in response to infection with *Phytophthora infestans*, associated with the accumulation of the isoflavonoid phytoalexin glyceollin (Bhattacharyya, M. K. and E. W. B. Ward, 1987, "Biosynthesis and metabolism of glyceollin I in soybean hypocotyls following wounding or inoculation with *Phytophthora megasperma* f. sp. *glycinea*," *Physiol Mol Plant Path* 31: 387-405). It was also known that an enzyme catalyzing a similar reaction to IFS, namely the 2-hydroxylation of flavanone but without aryl migration, belongs to the CYP93B1 subclass of cytochrome P450s (Akashi, et al, 1998, "Identification of a cytochrome P450 cDNA encoding (2S)-flavanone 2-hydroxylase of licorice (*Glycyrrhiza echinata* L.: *Fabaceae*) which represents licodione synthase and flavone synthase II," *FEBS Letters* 431: 287-290). We therefore searched an expressed sequence tag (EST) database of partial soybean sequences obtained by mass sequencing of two cDNA libraries: a *Phytophthora*-infected hypocotyl cDNA library (48 hours after infection) and a mid to late developmental stage seed library. Nine candidate P450 sequences were identified, of which three belonged to the CYP93 family. DNA probes were made from the EST clones of the three CYP93 candidates and were used to probe an RNA blot of transcripts from alfalfa suspension cells at various times after exposure to yeast elicitor, a treatment known to induce IFS activity at the onset of isoflavonoid phytoalexin accumulation (Kessmann, et al., 1990, "Stress responses in alfalfa (*Medicago sativa* L.) III. Induction of medicarpin and cytochrome P450 enzyme activities in elicitor-treated cell suspension cultures and protoplasts," *Plant Cell Reports* 9: 38-41). One P450 probe cross-hybridized and detected alfalfa transcripts that were strongly induced by elicitation. This probe was derived from a clone with high homology to soybean CYP93C1 as described below, and the insert in the EST clone was full length. The insert was excised and then cloned into the baculovirus expression system for functional identification by heterologous expression in insect cells (Pauli, H. H. and T. M. Kutchan, 1998, "Molecular cloning and functional heterologous expression of two alleles encoding (S)-*N*-methylcoclaurine 3'-

hydroxylase (CYP80B1), a new methyl jasmonate-inducible cytochrome P-450-dependent mono-oxygenase of benzyloquinoline alkaloid biosynthesis," *The Plant J* 13: 793-801).

5 The carbon monoxide difference spectrum of microsomes isolated from insect cells expressing the soybean CYP93C clone indicated the presence of expressed cytochrome P450, as seen from an absorption peak at 450 nm that was not present in similar spectra from insect microsomes originating from cells transformed with a control vector. Unlabeled liquiritigenin was then fed to the microsomes in the presence of NADPH. The substrate remained unconverted in microsomes from cells
10 harboring the control vector. However, in microsomes expressing the CYP93C clone, a new peak of RT 29.96 min was observed by high performance liquid chromatography (Fig 8A). The amount of this peak was reduced 10-fold if NADPH was omitted from the incubations (Fig. 8B). The UV spectrum of the product, obtained by diode array detection, was identical to that of authentic daidzein (λ_{max}
15 248 nm, sh 302 nm, λ_{min} 222 nm). The product was collected, derivatized, and analyzed by GC-MS. The mass spectrum of the BSTFA derivative was identical to that of an authentic sample of daidzein (Fig. 9). Microsomes containing the CYP93C clone also metabolized naringenin to yield genistein, although somewhat less efficiently than the reaction with liquiritigenin (Fig. 8C). Insect cell microsomes
20 expressing a different soybean cytochrome P450 cDNA, CYP93E, did not convert liquiritigenin to daidzein when incubated in the presence of NADPH (Fig. 8D). These results indicate that the soybean CYP93C encodes IFS.

Example 3: Method of Increasing Dietary Isoflavonoid Intake

Transgenic tomato plants are produced by the introduction of *CYP93C1v2* via
25 standard *Agrobacterium*-based procedures. In a preferred embodiment, the *CYP93C1v2* coding sequence is under control of a gene promoter giving specific expression in the fruit. Progeny containing the coding region of the *CYP93C1v2* gene are selected at the seedling stage by standard polymerase chain reaction and/or DNA blot analysis known to those skilled in the art. Plants scoring positive for possession
30 of the transgene are grown to fruiting, and fruit analyzed for the presence of

isoflavones by the HPLC methods shown in Fig. 7 and Fig. 8 of the present invention. Fruit harvested from the transgenic tomato plants are ingested to increase the dietary intake of isoflavonoids.

5 It is to be understood that the above description is of preferred exemplary embodiments of the invention and is intended to be illustrative of the invention, but is not to be construed to limit the scope of the invention in any way. Modifications may be made in the structural features of the invention without departing from the scope of the invention.

10 In summary, isoflavones can now be genetically engineered to provide potential human health benefits of dietary isoflavones and to increase disease resistance in plants. Isoflavones can now be produced in transgenic plants species in which isoflavones do not naturally occur, i.e., in species other than legumes. For example, engineering constitutive production of daidzein and/or genistein or their conjugates into tomato, potato, corn, or other popular components of the human diet,
15 leads to human health benefits, such as reduced cancer risk, reduced incidence of osteoporosis, and treatment for alcoholism. Alternatively, introducing infection-inducible isoflavonoid biosynthesis into non-legumes qualitatively complements these plants' phytoalexin defenses against microbial pathogens, whereas over-expression of the isoflavonoid pathway in legumes quantitatively increases this defense response.
20 Finally, modifying the extent of production of isoflavonoids in legume roots positively impacts nodulation efficiency and therefore plant yield.

WE CLAIM:

1. A method for introducing into a naturally non-isoflavonoid-producing plant species the enzyme catalyzing the aryl migration of a flavanone to form an isoflavanone intermediate or an isoflavone, comprising:
introducing a DNA segment encoding said enzyme into said plant to form a
5 transgenic plant, wherein said transgenic plant expresses said DNA segment under the control of a suitable constitutive or inducible promoter when said transgenic plant is exposed to conditions which permit expression.
2. The method of Claim 1, wherein said DNA segment comprises isolated genomic DNA.
3. The method of Claim 1, wherein said DNA segment comprises recombinant DNA.
4. The method of Claim 1, 2 or 3, wherein said DNA segment comprises a CYP93C gene.
5. The method of Claim 4, wherein said DNA segment consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.
6. The method of Claim 1, 2 or 3, wherein said DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene.
7. The method of Claim 6, wherein said DNA segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.
8. The method of Claim 1, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant to cause in vivo formation of daidzein or a daidzein derivative.

9. The method of Claim 2, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant to cause in vivo formation of daidzein or a daidzein derivative.
10. The method of Claim 3, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant to cause in vivo formation of daidzein or a daidzein derivative.
11. The method of Claim 8, 9 or 10, wherein said plant is further transformed to comprise said chalcone synthase, chalcone reductase, and chalcone isomerase genes.
12. The method of Claim 1, 2, 3, 8, 9 or 10, wherein said plant further comprises downstream genes to metabolize said formed isoflavanone intermediate or isoflavone to biologically active isoflavonoid derivatives or conjugates.
13. The method of Claim 12, wherein said downstream gene is selected from the group consisting of isoflavone *O*-methyltransferase, isoflavone 2'-hydroxylase, isoflavone reductase, and vestitone reductase.
14. The method of Claim 13, wherein said plant comprises downstream gene 4'-*O*-methyltransferase to form biochanin A or a biochanin A derivative.
15. The method of Claim 1, 2, 3, 5, 7, 8, 9, 10, 13 or 14, wherein said flavanone is liquiritigenin.
16. The method of Claim 4, wherein said flavanone is liquiritigenin.
17. The method of Claim 6, wherein said flavanone is liquiritigenin.
18. The method of Claim 11, wherein said flavanone is liquiritigenin.
19. The method of Claim 12, wherein said flavanone is liquiritigenin.
20. The method of Claim 1, 2, 3, 5, 7, 8, 9, 10, 13 or 14, wherein said flavanone is naringenin.

21. The method of Claim 4, wherein said flavanone is naringenin.
22. The method of Claim 6, wherein said flavanone is naringenin.
23. The method of Claim 11, wherein said flavanone is naringenin.
24. The method of Claim 12, wherein said flavanone is naringenin.
25. A method for increasing the level of isoflavonoid compounds in naturally isoflavonoid-producing plants comprising:
 - introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to yield an isoflavonoid to form a transgenic plant, wherein
 - 5 said transgenic plant expresses said DNA segment under the control of a suitable constitutive or inducible promoter when said transgenic plant is exposed to conditions which permit expression.
26. The method of Claim 25, wherein said isoflavonoid is selected from the group consisting of an isoflavanone intermediate, an isoflavone, an isoflavone derivative, and an isoflavone conjugate.
27. The method of Claim 25, wherein said DNA segment comprises isolated genomic DNA.
28. The method of Claim 26, wherein said DNA segment comprises isolated genomic DNA.
29. The method of Claim 25, wherein said DNA segment comprises recombinant DNA.
30. The method of Claim 26, wherein said DNA segment comprises recombinant DNA.
31. The method of Claim 25, 26, 27, 28, 29 or 30, wherein said DNA segment comprises a CYP93C gene.

32. The method of Claim 31, wherein said DNA segment consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

33. The method of Claim 25, 26, 27, 28, 29 or 30, wherein said DNA segment is a *Medicago truncatula* homolog of a CYP93C gene.

34. The method of Claim 33, wherein said DNA segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

35. The method of Claim 25, 26, 27, 28, 29, 30, 32 or 34, wherein said flavanone is liquiritigenin.

36. The method of Claim 31, wherein said flavanone is liquiritigenin.

37. The method of Claim 33, wherein said flavanone is liquiritigenin.

38. The method of Claim 25, 26, 27, 28, 29, 30, 32 or 34, wherein said flavanone is naringenin.

39. The method of Claim 31, wherein said flavanone is naringenin.

40. The method of Claim 33, wherein said flavanone is naringenin.

41. A method for synthesizing an isoflavanone intermediate or an isoflavone from a flavanone by expressing a recombinant CYP93C gene segment in a suitable bacterial, fungal, algal, or insect cell system.

42. The method of Claim 41, wherein said gene segment consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

43. The method of Claim 41, wherein said gene segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

44. A method of reducing the levels of isoflavonoid compounds in a naturally isoflavonoid-producing plant comprising introducing and expressing an antisense or gene silencing construct that contains an intact CYP93C gene or segments thereof into said plant.

45. The method of Claim 44, wherein said gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

46. The method of Claim 44, wherein said gene consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

47. A naturally non-isoflavonoid-producing plant cell transformed by introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to form an isoflavanone intermediate or an isoflavone, wherein said transgenic plant cell expresses said DNA segment under the control of a suitable
5 constitutive or inducible promoter when exposed to conditions which permit expression.

48. The plant cell of Claim 47, wherein said DNA segment comprises isolated genomic DNA.

49. The plant cell of Claim 47, wherein said DNA segment comprises recombinant DNA.

50. The plant cell of Claim 47, 48 or 49, wherein said DNA segment comprises a CYP93C gene.

51. The plant cell of Claim 50, wherein said DNA segment consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

52. The plant cell of Claim 47, 48 or 49, wherein said DNA segment comprises a *Medicago truncatula* homolog of a CYP93C gene.

53. The plant cell of Claim 52, wherein said DNA segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

54. The plant cell of Claim 47, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant cell to cause in vivo formation of daidzein or a daidzein derivative.

55. The plant cell of Claim 48, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant cell to cause in vivo formation of daidzein or a daidzein derivative.

56. The plant cell of Claim 49, wherein chalcone synthase, chalcone reductase, and chalcone isomerase genes are also expressed in said plant cell to cause in vivo formation of daidzein or a daidzein derivative.

57. The plant cell of Claim 54, 55 or 56, wherein said plant cell is further transformed to comprise said chalcone synthase, chalcone reductase, and chalcone isomerase genes.

58. The plant cell of Claim 47, 48, 49, 54, 55 or 56, wherein said plant cell further comprises downstream genes to metabolize said formed isoflavanone intermediate or isoflavone to biologically active isoflavonoid derivatives or conjugates.

59. The plant cell of Claim 58, wherein said downstream gene is selected from the group consisting of isoflavone *O*-methyltransferase, isoflavone 2'-hydroxylase, isoflavone reductase, and vestitone reductase.

60. The plant cell of Claim 59, wherein said plant cell comprises downstream gene 4'-*O*-methyltransferase to cause in vivo formation of biochanin A or a biochanin A derivative.

61. A naturally isoflavonoid-producing plant cell transformed by introducing a DNA segment encoding the enzyme catalyzing the aryl migration of a flavanone to yield an isoflavonoid to form a transformed plant cell, wherein said transformed plant cell expresses said DNA segment under the control of a suitable constitutive or
5 inducible promoter when exposed to conditions which permit expression.

62. The plant cell of Claim 61, wherein said isoflavonoid is selected from the group consisting of an isoflavanone intermediate, an isoflavone, an isoflavone derivative, and an isoflavone conjugate.

63. The plant cell of Claim 61, wherein said DNA segment comprises isolated genomic DNA.

64. The plant cell of Claim 62, wherein said DNA segment comprises isolated genomic DNA.

65. The plant cell of Claim 61, wherein said DNA segment comprises recombinant DNA.

66. The plant cell of Claim 62, wherein said DNA segment comprises recombinant DNA.

67. The plant cell of Claim 61, 62, 63, 64, 65 or 66, wherein said DNA segment comprises a CYP93C gene.

68. The plant cell of Claim 67, wherein said DNA segment consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

69. The plant cell of Claim 61, 62, 63, 64, 65 or 66, wherein said DNA segment is a *Medicago truncatula* homolog of a CYP93C gene.

70. The plant cell of Claim 69, wherein said DNA segment consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

71. A transgenic plant cell having reduced levels of isoflavonoid compounds, said plant cell transformed by introducing an antisense or gene silencing construct that contains an intact CYP93C gene or segments thereof into said plant cell.

72. The plant cell of Claim 71, wherein said gene consists essentially of the sequence from about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

73. The plant cell of Claim 71, wherein said gene consists essentially of the sequence from about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4.

74. An isolated gene or DNA segment comprising a portion which encodes a cytochrome P450 of the CYP93 family that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said portion consists essentially of about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

75. The gene or DNA segment of Claim 74, wherein said gene is the soybean gene encoding the enzyme catalyzing the aryl migration of liquiritigenin.

76. The gene or DNA segment of Claim 74, wherein said gene is the soybean gene encoding the enzyme catalyzing the aryl migration of naringenin.

77. A protein encoded by a portion of an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said portion consists essentially of about nucleotide 36 to about nucleotide 1598 of the sequence depicted in SEQ ID NO:1.

78. An isolated gene or DNA segment comprising a portion which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said portion is a *Medicago truncatula* homolog of a CYP93C gene.

79. The gene or DNA segment of Claim 78 consisting essentially of about nucleotide 92 to about nucleotide 1657 of the sequence depicted in SEQ ID NO:4

80. The gene or DNA segment of Claim 78, wherein said gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of liquiritigenin.

81. The gene or DNA segment of Claim 79, wherein said gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of liquiritigenin.

82. The gene or DNA segment of Claim 78, wherein said gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of naringenin.

83. The gene or DNA segment of Claim 79, wherein said gene is the *Medicago truncatula* gene encoding the enzyme catalyzing the aryl migration of naringenin.

84. A protein encoded by a portion of an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said portion is a *Medicago truncatula* homolog of a CYP93C gene.

85. A food comprising edible transgenic plant material capable of being ingested for its nutritional value, said transgenic plant is transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone,
5 wherein said transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

86. A food comprising at least one isoflavonoid, wherein said isoflavonoid is isolated from a transgenic plant transformed with an isolated gene or DNA segment

which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant
5 exhibits increased levels of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

87. A composition comprising at least a portion of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits increased levels
5 of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment, said composition suitable for ingestion as a food stuff, a nutritional supplement, an animal feed supplement, or a nutraceutical.

88. A composition comprising an isoflavonoid suitable for administration as a food stuff, a nutritional supplement, an animal feed supplement, a nutraceutical, or a pharmaceutical, said isoflavonoid isolated from at least a portion of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450
5 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits increased levels of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

89. A method of increasing the nutritional value of a plant by transforming said plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits increased levels
5 of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

90. A method of using a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment, to provide a nutraceutical benefit to a human or animal administered said isoflavonoid.

91. The method of claim 90, wherein said isoflavonoid is administered by ingestion of at least a portion of said plant.

92. The method of claim 90, wherein said isoflavonoid is administered by ingestion of a composition comprising an isoflavonoid isolated from said plant.

93. A method of using an isoflavonoid isolated from a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment, to provide a pharmaceutical benefit to a patient administered said isoflavonoid.

94. A method of increasing disease resistance in a plant by transforming said plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

95. A method of increasing nodulation efficiency of a leguminous plant by transforming said plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid released from the roots of said transgenic plant when compared to the level of the isoflavonoid released from the roots of plants of the same species which do not comprise said isolated gene or DNA segment.

96. A leguminous transgenic plant exhibiting increased nodulation efficiency, wherein said transgenic plant is transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, and wherein said transgenic plant exhibits an increased level of an isoflavonoid released from the roots of said transgenic plant when compared to the level of the isoflavonoid released from the roots of plants of the same species which do not comprise said isolated gene or DNA segment.

97. A method of increasing bacterial or fungal symbiosis in a plant by transforming said plant with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment.

98. A transgenic plant exhibiting an increased level of bacterial or fungal symbiosis, wherein said transgenic plant is transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, and wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise the isolated gene or DNA segment.

99. A transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said recombinant DNA sequence.

100. Seed from a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said recombinant DNA sequence.

101. Progeny from a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said recombinant DNA sequence.

102. Progeny from seed of a transgenic plant comprising at least one recombinant DNA sequence encoding a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said recombinant DNA sequence.

103. Use of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said

isolated gene or DNA segment, for the preparation of a nutraceutical preparation for achieving a nutritional effect.

104. Use of a transgenic plant transformed with an isolated gene or DNA segment which encodes a cytochrome P450 that can catalyze the aryl migration of a flavanone to yield an isoflavanone intermediate or an isoflavone, wherein said transgenic plant exhibits an increased level of an isoflavonoid when compared to the level of said isoflavonoid in plants of the same species which do not comprise said isolated gene or DNA segment, for the preparation of a pharmaceutical preparation for achieving a therapeutic effect.
- 5

1/15




Fig. 1

2/15

```

1   GAGCAAAGAT CAAACAAACC AAGGACGAGA ACACGATGTT GCTTGAACCTT
51  GCACTTGGTT TATTGGTTTT GGCTCTGTTT CTGCACTTGC GTCCCAACACC
101 CACTGCAAAA TCAAAAGCAC TTCGCCATCT CCCAAACCCA CCAAGCCCAA
151 AGCCTCGTCT TCCCTTCATA GGACACCTTC ATCTCTTAAA AGACAAACTT
201 CTCCACTACG CACTCATCGA CCTCTCCAAA AAACATGGTC CCTTATCTCTC
251 TCTCTACTTT GGCTCCATGC CAACCGTTGT TGCCTCCACA CCAGAATTGT
301 TCAAGCTCTT CCTCCAAACG CACGAGGCAA CTTCTTCAA CACAAGGTTT
351 CAAACCTCAG CCATAAGACG CCTCACCTAT GATAGCTCAG TGGCCATGGT
401 TCCCTTCGGA CCTTACTGGA AGTTCGTGAG GAAGCTCATC ATGAACGACC
451 TTCTCAACGC CACCACTGTA AACAAGTTGA GGCCTTTGAG GACCCAACAG
501 ATCCGCAAGT TCCTTAGGGT TATGGCCCAA GGCGCAGAGG CACAGAAGCC
551 CCTTGACTTG ACCGAGGAGC TTCTGAAATG GACCAACAGC ACCATCTCCA
601 TGATGATGCT CGGCGAGGCT GAGGAGATCA GAGACATCGC TCGCGAGGTT
651 CTTAAGATCT TTGGCGAATA CAGCCTCACT GACTTCATCT GGCCATTGAA
701 GCATCTCAAG GTTGGAAAGT ATGAGAAGAG GATCGACGAC ATCTTGAACA
751 AGTTCGACCC TGTCGTTGAA AGGGTCATCA AGAAGCGCCG TGAGATCGTG
801 AGGAGGAGAA AGAACGGAGA GGTGTTGAG GGTGAGGTCA GCGGGGTTTT
851 CCTTGACACT TTGCTTGAAT TCGCTGAGGA TGAGACCATG GAGATCAAAA
901 TCACCAAGGA CCACATCAAG GGTCTTGTTG TCGACTTTTT CTCCGCAGGA
951 ACAGACTCCA CAGCGGTGGC AACAGAGTGG GCATTGGCAG AACTCATCAA
1001 CAATCCTAAG GTGTTGGAAA AGGCTCGTGA GGAGGTCTAC AGTGTGTGG
1051 GAAAGGACAG ACTTGTGGAC GAAAGTTGACA CTCAAAACCT TCCTTACATT
1101 AGAGCAATCG TGAAGGAGAC ATTCCGCATG CACCCGCCAC TCCAGTGGT
1151 CAAAAGAAAG TGCACAGAAG AGTGTGAGAT TAATGGATAT GTGATCCAG
1201 AGGGAGCATT GATTCTCTTC AATGTATGGC AAGTAGGAAG AGACCCCAA
1251 TACTGGGACA GACCATCGGA GTTCCGTCCT GAGAGGTTC TAGAGACAGG
1301 GGCTGAAGGG GAAGCAGGGC CTCTTGATCT TAGGGGACAA CATTTC AAC
1351 TTCTCCCAT TGGGTCTGGG AGGAGAATGT GCCCTGGAGT CAATCTGGCT
1401 ACTTCGGGAA TGGCAACACT TCTTGCATCT CTTATTAGT GCTTCGACTT
1451 GCAAGTGCTG GGTCCACAAG GACAGATATT GAAGGGTGGT GACGCCAAAG
1501 TTAGCATGGA AGAGAGAGCC GGCCTCACTG TTCCAAGGGC ACATAGTCTT
1551 GTCTGTGTTC CACTTGCAAG GATCGGCGTT GCATCTAAAC TCCTTTCTTA
1601 ATTAAGATCA TCGTCATCAT CATCATATGT AATATTTACT TTTTGTGTGT
1651 TGATAATCAT CATTTCAATA AGGTCTCATT CATCTACTTT TTATGAAGTA
1701 TATAAGCCCT TCCATGC

```

Fig. 2

3/15

[illegible]

Fig. 3

4/15

```

1   CAACACCTAA GAGTAACTAA TAAGAACTTT CTTTCTACTT CTTAGTATAC
51  TTAACAACTT AAGTAAATAT ACTACAAAGA AGCTATACAC CATGTTGGTG
101 GAACCTGCAG TTACTCTATT GCTCATTGCT CTCTTCTTAC ACTTGCGTCC
151 AACACCTACT GCAAAATCAA AGGCTCTTCG CCACCTTCCA AATCCACCAA
201 GCCCTAAACC ACGTCTTCCA TTCATAGGTC ATCTTCACCT TTTGGATAAC
251 CCACTTCTTC ACCACACTCT TATCAAGTTA GGAAAGCGTT ATGGCCCTTT
301 GTACACTCTT TACTTTGGTT CCATGCCTAC CGTTGTTGCA TCCACTCCTG
351 ACTTGTTTAA ACTTTTCCTT CAAACCCATG AAGCTACTTC CTTTAACACA
401 AGATTCCAAA CCTCTGCTAT TAGTCGTCTT ACCTATGACA ACTCTGTTGC
451 TATGGTTCCA TTTGCACCTT ATTGGAAGTT TATTAGAAAG CTTATCATGA
501 ACGACTTGCT CAACGCCACC ACTGTTAACA AATTGAGGCC ATTGAGGAGC
551 CGAGAAATCC TTAAGGTCTT TAAGGTCATG GCTAATAGTG CTGAAACTCA
601 ACAGCCACTT GATGTCAC TG AGGAGCTTCT CAAGTGGACA AACAGCACAA
651 TCTCTACCAT GATGTTGGGT GAGGCCGAAG AGGTTAGAGA TATTGCTCGT
701 GATGTTCTTA AGATCTTTGG AGAATATAGT GTTACAAACT TTATTTGGCC
751 TTTGAACAAG TTTAAGTTTG GAAACTATGA TAAGAGAACT GAGGAGATTT
801 TCAATAAGTA TGATCCTATC ATTGAAAAGG TTATCAAGAA ACGACAAGAG
851 ATTGTAACA AAAGAAAAAA TGGAGAAATC GTAGAAGGCG AGCAGAATGT
901 TGTTTTCTT GACACTTTCG TTGAATTGCA ACAAGATGAG ACCATGGAGA
951 TCAAAATTAC AAAGGAACAA ATCAAGGGTC TTGTTGTGGA TTTTTTCTCT
1001 GCAGGAACAG ACTCCACCGC CGTGTCTACA GAATGGACTT TATCAGAGCT
1051 CATCAATAAT CCTAGAGTGT TGAAGAAAGC TCGAGAGGAG ATTGACTCTG
1101 TTGTGGGAAA AGATAGACTG GTTGATGAAT CAGATGTTCA GAATCTTCCT
1151 TACATTAAAG CCATCGTAAA AGAAGCATTT CGCTTGCACC CACCCTACC
1201 TGTAGTCAAA AGAAAATGTA CACAAGAATG TGAGATCGAC GGGTATGTGG
1251 TTCCAGAAGG AGCACTAATA CTTTCAATG TCTGGGCAGT GGAAGAGAC
1301 CCAAAATATT GGGTAAAGCC ATTGGAATTT CGTCCAGAGA GGTTCATAGA
1351 AAATGTTGGT GAAGGTGAAG CAGCTTCAAT TGATCTTAGG GGTCAACATT
1401 TCACACTTCT ACCATTTGGG TCTGGAAGAA GGATGTGTCC TGGAGTCAAT
1451 TTGGCTACTG CAGGAATGGC CACAATGATT GCATCTATTA TCCAATGCCT
1501 CGATCTCCAA GTACCTGGTC AACATGGAGA AATATTGAAT GGTGATTATG
1551 CTAAGGTTAG CATGGAAGAG AGACCTGGTC TCACAGTTCC AAGGGCACAT
1601 AATCTCATGT GTGTTCTCTT TGCAAGAGCT GGTGTCGCAG ATAACTTCT
1651 TTCCTCTAA AATATCTTGA GAGGATGAAT CACCAACATA TAGCCTCTCT
1701 TTGGTACTAC AAAATTATGA TGTAATTTTC TTATTTTTTC TGTCACAAAG
1751 GAAGTGTTGT AACTTGTAAT TGCATACAAA ATCTATAAAT TTTATCATCC
1801 TATTCATTAT T

```

Fig. 4

5/15

```

mtIFS      1  MLVELAVTLLLLIALFLHLRPTPTAKSKALRHLPNPPSPKPRLPFFIGHLHL 50
           ||.|||. ||.:|||||||||||||||||||||||||||||||||
CYP93C1v2  1  MLELALGLLVLALFLHLRPTPTAKSKALRHLPNPPSPKPRLPFFIGHLHL 50
           .
51  LDNPLLHHTLIKLGKRYGPLYTLYFGSMPTVVASTPDLFKLFLQTHEATS 100
           | . |||: || | |.:|||.|||||||||||||:|||||||
51  LKDKLLHYALIDLKSKHGPLFSLYFGSMPTVVASTPELFLKLFLQTHEATS 100
           .
101 FNTRFQTSAISRLTYDNSVAMVPFAPYWKFIRKLIMNDLLNATTVNKLRLP 150
           ||||||| . ||||. ||||| |||||:|||||||
101 FNTRFQTSAIRRLTYDSSVAMVPFGPYWKFVRKLIMNDLLNATTVNKLRLP 150
           .
151 LRSREILKVLKVMANSAETQOPLDVTEELLKWTNSTISTMMLGEAEVVD 200
           ||.:| | |.:||| | |.|||.||||||| |||||:|
151 LRTQQIRKFLRVMAQGAQAQKPLDLTEELLKWTNSTISMMLGEAEIIRD 200
           .
201 IARDVLKIFGEYSVTNFIWPLNKFKEGNYDKRTEEIFNKYDPIIEKVIKK 250
           |||:|||||||.||.|||| | | |.:|||:| |||:|:|
201 IAREVLKIFGEYSLTDFIWPLKHLKVGKYEKRIDDILNKFDPVVERVIKK 250
           .
251 RQEIVNKRKNGEIVEGEQNVVFLDTLLEFAQDETMEIKITKEQIKGLVVD 300
           |.||| :|||||:|||| . |||||||:|||||||: |||||
251 RREIVRRRKNGEVVEGEVSGVFLDTLLEFAEDETMEIKITKDHIKGLVVD 300
           .
301 FFSAGTDSTAVSTEWTLSELINNPRVLKKAREEIDSVVGKDRLVDESVDQ 350
           |||||||||.||| |.|||||:||.|||||: ||||||||| |
301 FFSAGTDSTAVATEWALAEELINNPVKLEKAREEVSVVGKDRLVDEVDQ 350
           .
351 NLPYIKAIVKEAFRLHPPLPVVVRKCTQECEIDGYVVPFEGALILFNWVAV 400
           |||||:||||| ||:|||||||||||:||||.|||:||||||| |
351 NLPYIRAIVKETFRMHPPLPVVVRKCTEECEINGYVIPEGALILFNWVQV 400
           .
401 GRDPKYWVKPLEFRPERFIENVGEGEAASIDLRGQHFTLLPFGSRRMCP 450
           ||||||| :| |||||||:| |||| :|||||| | |||||||
401 GRDPKYWDRPSEFRPERFLETGAEGEAGPLDLRGQHFQLLPFGSRRMCP 450
           .
451 GVNLATAGMATMIASIIQCFDLQVPGQHGEILNGDYAKVSMEERPGLTVP 500
           ||||||.|||||:||:||||||| | |.:|| | ||||||| ||||
451 GVNLATSGMATLLASLIQCFDLQVLGPQGQILKGGDAKVSMEERAGLTVP 500
           .
501 RAHNLMCVPLARAGVADKLLSS 522
           |||.||.||||| ||| ||||
501 RAHSLVCVPLARIGVASKLLS 521
           .

```

Fig. 5

6/15




Fig. 6A

7/15




Fig. 6B




Fig. 6C

9/15




Fig. 7A

10/15




Fig. 7B

11/15




Fig. 7C

12/15




Fig. 7D




Fig. 8A




Fig. 8B

14/15




Fig. 8C






Fig. 8D

15/15

Fig. 9A**Fig. 9B**

SEQUENCE LISTING

<110> THE SAMUEL ROBERTS NOBLE FOUNDATION, INC.
 STEELE, Christopher L.
 DIXON, Richard A.

<120> GENETIC MANIPULATION OF ISOFLAVONOIDS

<130> 11137/05002

<140>

<141>

<150> 60/123,267

<151> 1999-03-08

<160> 5

<170> PatentIn Ver. 2.1

<210> 1

<211> 1717

<212> DNA

<213> Glycine max

<220>

<221> CDS

<222> (36)..(1598)

<400> 1

```

gagcaaagat caaacaacc aaggacgaga acacg atg ttg ctt gaa ctt gca      53
                                   Met Leu Leu Glu Leu Ala
                                   1           5

ctt ggt tta ttg gtt ttg gct ctg ttt ctg cac ttg cgt ccc aca ccc      101
Leu Gly Leu Leu Val Leu Ala Leu Phe Leu His Leu Arg Pro Thr Pro
              10              15              20

act gca aaa tca aaa gca ctt cgc cat ctc cca aac cca cca agc cca      149
Thr Ala Lys Ser Lys Ala Leu Arg His Leu Pro Asn Pro Pro Ser Pro
              25              30              35

aag cct cgt ctt ccc ttc ata gga cac ctt cat ctc tta aaa gac aaa      197
Lys Pro Arg Leu Pro Phe Ile Gly His Leu His Leu Leu Lys Asp Lys
              40              45              50

ctt ctc cac tac gca ctc atc gac ctc tcc aaa aaa cat ggt ccc tta      245
Leu Leu His Tyr Ala Leu Ile Asp Leu Ser Lys Lys His Gly Pro Leu
              55              60              65              70

ttc tct ctc tac ttt ggc tcc atg cca acc gtt gtt gcc tcc aca cca      293
Phe Ser Leu Tyr Phe Gly Ser Met Pro Thr Val Val Ala Ser Thr Pro
              75              80              85

gaa ttg ttc aag ctc ttc ctc caa acg cac gag gca act tcc ttc aac      341
Glu Leu Phe Lys Leu Phe Leu Gln Thr His Glu Ala Thr Ser Phe Asn
              90              95              100

```

aca agg ttc caa acc tca gcc ata aga cgc ctc acc tat gat agc tca	389
Thr Arg Phe Gln Thr Ser Ala Ile Arg Arg Leu Thr Tyr Asp Ser Ser	
105 110 115	
gtg gcc atg gtt ccc ttc gga cct tac tgg aag ttc gtg agg aag ctc	437
Val Ala Met Val Pro Phe Gly Pro Tyr Trp Lys Phe Val Arg Lys Leu	
120 125 130	
atc atg aac gac ctt ctc aac gcc acc act gta aac aag ttg agg cct	485
Ile Met Asn Asp Leu Leu Asn Ala Thr Thr Val Asn Lys Leu Arg Pro	
135 140 145 150	
ttg agg acc caa cag atc cgc aag ttc ctt agg gtt atg gcc caa ggc	533
Leu Arg Thr Gln Ile Arg Lys Phe Leu Arg Val Met Ala Gln Gly	
155 160 165	
gca gag gca cag aag ccc ctt gac ttg acc gag gag ctt ctg aaa tgg	581
Ala Glu Ala Gln Lys Pro Leu Asp Leu Thr Glu Glu Leu Leu Lys Trp	
170 175 180	
acc aac agc acc atc tcc atg atg atg ctc ggc gag gct gag gag atc	629
Thr Asn Ser Thr Ile Ser Met Met Met Leu Gly Glu Ala Glu Glu Ile	
185 190 195	
aga gac atc gct cgc gag gtt ctt aag atc ttt ggc gaa tac agc ctc	677
Arg Asp Ile Ala Arg Glu Val Leu Lys Ile Phe Gly Glu Tyr Ser Leu	
200 205 210	
act gac ttc atc tgg cca ttg aag cat ctc aag gtt gga aag tat gag	725
Thr Asp Phe Ile Trp Pro Leu Lys His Leu Lys Val Gly Lys Tyr Glu	
215 220 225 230	
aag agg atc gac gac atc ttg aac aag ttc gac cct gtc gtt gaa agg	773
Lys Arg Ile Asp Asp Ile Leu Asn Lys Phe Asp Pro Val Val Glu Arg	
235 240 245	
gtc atc aag aag cgc cgt gag atc gtg agg agg aga aag aac gga gag	821
Val Ile Lys Lys Arg Arg Glu Ile Val Arg Arg Arg Lys Asn Gly Glu	
250 255 260	
gtt gtt gag ggt gag gtc agc ggg gtt ttc ctt gac act ttg ctt gaa	869
Val Val Glu Gly Glu Val Ser Gly Val Phe Leu Asp Thr Leu Leu Glu	
265 270 275	
ttc gct gag gat gag acc atg gag atc aaa atc acc aag gac cac atc	917
Phe Ala Glu Asp Glu Thr Met Glu Ile Lys Ile Thr Lys Asp His Ile	
280 285 290	
aag ggt ctt gtt gtc gac ttt ttc tcg gca gga aca gac tcc aca gcg	965
Lys Gly Leu Val Val Asp Phe Phe Ser Ala Gly Thr Asp Ser Thr Ala	
295 300 305 310	
gtg gca aca gag tgg gca ttg gca gaa ctc atc aac aat cct aag gtg	1013
Val Ala Thr Glu Trp Ala Leu Ala Glu Leu Ile Asn Asn Pro Lys Val	
315 320 325	
ttg gaa aag gct cgt gag gag gtc tac agt gtt gtg gga aag gac aga	1061
Leu Glu Lys Ala Arg Glu Glu Val Tyr Ser Val Val Gly Lys Asp Arg	
330 335 340	

```

ctt gtg gac gaa gtt gac act caa aac ctt cct tac att aga gca atc 1109
Leu Val Asp Glu Val Asp Thr Gln Asn Leu Pro Tyr Ile Arg Ala Ile
345 350 355

gtg aag gag aca ttc cgc atg cac ccg cca ctc cca gtg gtc aaa aga 1157
Val Lys Glu Thr Phe Arg Met His Pro Pro Leu Pro Val Val Lys Arg
360 365 370

aag tgc aca gaa gag tgt gag att aat gga tat gtg atc cca gag gga 1205
Lys Cys Thr Glu Glu Cys Glu Ile Asn Gly Tyr Val Ile Pro Glu Gly
375 380 385 390

gca ttg att ctc ttc aat gta tgg caa gta gga aga gac ccc aaa tac 1253
Ala Leu Ile Leu Phe Asn Val Trp Gln Val Gly Arg Asp Pro Lys Tyr
395 400 405

tgg gac aga cca tcg gag ttc cgt cct gag agg ttc cta gag aca ggg 1301
Trp Asp Arg Pro Ser Glu Phe Arg Pro Glu Arg Phe Leu Glu Thr Gly
410 415 420

gct gaa ggg gaa gca ggg cct ctt gat ctt agg gga caa cat ttt caa 1349
Ala Glu Gly Glu Ala Gly Pro Leu Asp Leu Arg Gly Gln His Phe Gln
425 430 435

ctt ctc cca ttt ggg tct ggg agg aga atg tgc cct gga gtc aat ctg 1397
Leu Leu Pro Phe Gly Ser Gly Arg Arg Met Cys Pro Gly Val Asn Leu
440 445 450

gct act tcg gga atg gca aca ctt ctt gca tct ctt att cag tgc ttc 1445
Ala Thr Ser Gly Met Ala Thr Leu Leu Ala Ser Leu Ile Gln Cys Phe
455 460 465 470

gac ttg caa gtg ctg ggt cca caa gga cag ata ttg aag ggt ggt gac 1493
Asp Leu Gln Val Leu Gly Pro Gln Gly Gln Ile Leu Lys Gly Gly Asp
475 480 485

gcc aaa gtt agc atg gaa gag aga gcc ggc ctc act gtt cca agg gca 1541
Ala Lys Val Ser Met Glu Glu Arg Ala Gly Leu Thr Val Pro Arg Ala
490 495 500

cat agt ctt gtc tgt gtt cca ctt gca agg atc ggc gtt gca tct aaa 1589
His Ser Leu Val Cys Val Pro Leu Ala Arg Ile Gly Val Ala Ser Lys
505 510 515

ctc ctt tct taattaagat catcgatcatc atcatcatat gtaatatatta 1638
Leu Leu Ser
520

ctttttgtgt gttgataatc atcatttcaa taaggctctca ttcattctact ttttatgaag 1698
tatataagcc cttccatgc 1717

```

<210> 2

<211> 521

<212> PRT

<213> Glycine max

<400> 2

```

Met Leu Leu Glu Leu Ala Leu Gly Leu Leu Val Leu Ala Leu Phe Leu
 1           5           10           15

His Leu Arg Pro Thr Pro Thr Ala Lys Ser Lys Ala Leu Arg His Leu
          20           25           30

Pro Asn Pro Pro Ser Pro Lys Pro Arg Leu Pro Phe Ile Gly His Leu
          35           40           45

His Leu Leu Lys Asp Lys Leu Leu His Tyr Ala Leu Ile Asp Leu Ser
          50           55           60

Lys Lys His Gly Pro Leu Phe Ser Leu Tyr Phe Gly Ser Met Pro Thr
          65           70           75           80

Val Val Ala Ser Thr Pro Glu Leu Phe Lys Leu Phe Leu Gln Thr His
          85           90           95

Glu Ala Thr Ser Phe Asn Thr Arg Phe Gln Thr Ser Ala Ile Arg Arg
          100          105          110

Leu Thr Tyr Asp Ser Ser Val Ala Met Val Pro Phe Gly Pro Tyr Trp
          115          120          125

Lys Phe Val Arg Lys Leu Ile Met Asn Asp Leu Leu Asn Ala Thr Thr
          130          135          140

Val Asn Lys Leu Arg Pro Leu Arg Thr Gln Gln Ile Arg Lys Phe Leu
          145          150          155          160

Arg Val Met Ala Gln Gly Ala Glu Ala Gln Lys Pro Leu Asp Leu Thr
          165          170          175

Glu Glu Leu Leu Lys Trp Thr Asn Ser Thr Ile Ser Met Met Met Leu
          180          185          190

Gly Glu Ala Glu Glu Ile Arg Asp Ile Ala Arg Glu Val Leu Lys Ile
          195          200          205

Phe Gly Glu Tyr Ser Leu Thr Asp Phe Ile Trp Pro Leu Lys His Leu
          210          215          220

Lys Val Gly Lys Tyr Glu Lys Arg Ile Asp Asp Ile Leu Asn Lys Phe
          225          230          235          240

Asp Pro Val Val Glu Arg Val Ile Lys Lys Arg Arg Glu Ile Val Arg
          245          250          255

Arg Arg Lys Asn Gly Glu Val Val Glu Gly Glu Val Ser Gly Val Phe
          260          265          270

Leu Asp Thr Leu Leu Glu Phe Ala Glu Asp Glu Thr Met Glu Ile Lys
          275          280          285

Ile Thr Lys Asp His Ile Lys Gly Leu Val Val Asp Phe Phe Ser Ala
          290          295          300

Gly Thr Asp Ser Thr Ala Val Ala Thr Glu Trp Ala Leu Ala Glu Leu
          305          310          315          320

```

Ile Asn Asn Pro Lys Val Leu Glu Lys Ala Arg Glu Glu Val Tyr Ser
 325 330 335
 Val Val Gly Lys Asp Arg Leu Val Asp Glu Val Asp Thr Gln Asn Leu
 340 345 350
 Pro Tyr Ile Arg Ala Ile Val Lys Glu Thr Phe Arg Met His Pro Pro
 355 360 365
 Leu Pro Val Val Lys Arg Lys Cys Thr Glu Glu Cys Glu Ile Asn Gly
 370 375 380
 Tyr Val Ile Pro Glu Gly Ala Leu Ile Leu Phe Asn Val Trp Gln Val
 385 390 395 400
 Gly Arg Asp Pro Lys Tyr Trp Asp Arg Pro Ser Glu Phe Arg Pro Glu
 405 410 415
 Arg Phe Leu Glu Thr Gly Ala Glu Gly Glu Ala Gly Pro Leu Asp Leu
 420 425 430
 Arg Gly Gln His Phe Gln Leu Leu Pro Phe Gly Ser Gly Arg Arg Met
 435 440 445
 Cys Pro Gly Val Asn Leu Ala Thr Ser Gly Met Ala Thr Leu Leu Ala
 450 455 460
 Ser Leu Ile Gln Cys Phe Asp Leu Gln Val Leu Gly Pro Gln Gly Gln
 465 470 475 480
 Ile Leu Lys Gly Gly Asp Ala Lys Val Ser Met Glu Glu Arg Ala Gly
 485 490 495
 Leu Thr Val Pro Arg Ala His Ser Leu Val Cys Val Pro Leu Ala Arg
 500 505 510
 Ile Gly Val Ala Ser Lys Leu Leu Ser
 515 520

<210> 3

<211> 523

<212> PRT

<213> Glycyrrhiza echinata

<400> 3

Met Glu Pro Gln Leu Val Ala Val Ser Val Leu Val Ser Ala Leu Ile
 1 5 10 15
 Cys Tyr Phe Phe Phe Arg Pro Tyr Phe His Arg Tyr Gly Lys Asn Leu
 20 25 30
 Pro Pro Ser Pro Phe Phe Arg Leu Pro Ile Ile Gly His Met His Met
 35 40 45
 Leu Gly Pro Leu Leu His Gln Ser Phe His Asn Leu Ser His Arg Tyr
 50 55 60

Gly Pro Leu Phe Ser Leu Asn Phe Gly Ser Val Leu Cys Val Val Ala
 65 70 75 80
 Ser Thr Pro His Phe Ala Lys Gln Leu Leu Gln Thr Asn Glu Leu Ala
 85 90 95
 Phe Asn Cys Arg Ile Glu Ser Thr Ala Val Lys Lys Leu Thr Tyr Glu
 100 105 110
 Ser Ser Leu Ala Phe Ala Pro Tyr Gly Asp Tyr Trp Arg Phe Ile Lys
 115 120 125
 Lys Leu Ser Met Asn Glu Leu Leu Gly Ser Arg Ser Ile Asn Asn Phe
 130 135 140
 Gln His Leu Arg Ala Gln Glu Thr His Gln Leu Leu Arg Leu Leu Ser
 145 150 155 160
 Asn Arg Ala Arg Ala Phe Glu Ala Val Asn Ile Thr Glu Glu Leu Leu
 165 170 175
 Lys Leu Thr Asn Asn Val Ile Ser Ile Met Met Val Gly Glu Ala Glu
 180 185 190
 Glu Ala Arg Asp Val Val Arg Asp Val Thr Glu Ile Phe Gly Glu Phe
 195 200 205
 Asn Val Ser Asp Phe Ile Trp Leu Phe Lys Lys Met Asp Leu Gln Gly
 210 215 220
 Phe Gly Lys Arg Ile Glu Asp Leu Phe Gln Arg Phe Asp Thr Leu Val
 225 230 235 240
 Glu Arg Ile Ile Ser Lys Arg Glu Gln Thr Arg Lys Asp Arg Arg Arg
 245 250 255
 Asn Gly Lys Lys Gly Glu Gln Gly Ser Gly Asp Gly Ile Arg Asp Phe
 260 265 270
 Leu Asp Ile Leu Leu Asp Cys Thr Glu Asp Glu Asn Ser Glu Ile Lys
 275 280 285
 Ile Gln Arg Val His Ile Lys Ala Leu Ile Met Asp Phe Phe Thr Ala
 290 295 300
 Gly Thr Asp Thr Thr Ala Ile Ser Thr Glu Trp Ala Leu Val Glu Leu
 305 310 315 320
 Val Lys Lys Pro Ser Val Leu Gln Lys Val Arg Glu Glu Ile Asp Asn
 325 330 335
 Val Val Gly Lys Asp Arg Leu Val Glu Glu Ser Asp Cys Pro Asn Leu
 340 345 350
 Pro Tyr Leu Gln Ala Ile Leu Lys Glu Thr Phe Arg Leu His Pro Pro
 355 360 365
 Val Pro Met Val Thr Arg Arg Cys Val Ala Glu Cys Thr Val Glu Asn
 370 375 380

Tyr Val Ile Pro Glu Asp Ser Leu Leu Phe Val Asn Val Trp Ser Ile
 385 390 395 400
 Gly Arg Asn Pro Lys Phe Trp Asp Asn Pro Leu Glu Phe Arg Pro Glu
 405 410 415
 Arg Phe Leu Lys Leu Glu Gly Asp Ser Ser Gly Val Val Asp Val Arg
 420 425 430
 Gly Ser His Phe Gln Leu Leu Pro Phe Gly Ser Gly Arg Arg Met Cys
 435 440 445
 Pro Gly Val Ser Leu Ala Met Gln Glu Val Pro Ala Leu Leu Gly Ala
 450 455 460
 Ile Ile Gln Cys Phe Asp Phe His Val Val Gly Pro Lys Gly Glu Ile
 465 470 475 480
 Leu Lys Gly Asp Asp Ile Val Ile Asn Val Asp Glu Arg Pro Gly Leu
 485 490 495
 Thr Ala Pro Arg Ala His Asn Leu Val Cys Val Pro Val Asp Arg Thr
 500 505 510
 Ser Gly Gly Gly Pro Leu Lys Ile Ile Glu Cys
 515 520

<210> 4
 <211> 1811
 <212> DNA
 <213> *Medicago truncatula*

<220>
 <221> CDS
 <222> (92)..(1657)

<400> 4
 caacacctaagagtaactaa taagaacttt ctttctactt cttagtatac ttaacaactt 60
 aagtaaatat actacaaaga agctatacac c atg ttg gtg gaa ctt gca gtt 112
 Met Leu Val Glu Leu Ala Val
 1 5
 act cta ttg ctc att gct ctc ttc tta cac ttg cgt cca aca cct act 160
 Thr Leu Leu Leu Ile Ala Leu Phe Leu His Leu Arg Pro Thr Pro Thr
 10 15 20
 gca aaa tca aag gct ctt cgc cac ctt cca aat cca cca agc cct aaa 208
 Ala Lys Ser Lys Ala Leu Arg His Leu Pro Asn Pro Pro Ser Pro Lys
 25 30 35
 cca cgt ctt cca ttc ata ggt cat ctt cac ctt ttg gat aac cca ctt 256
 Pro Arg Leu Pro Phe Ile Gly His Leu His Leu Leu Asp Asn Pro Leu
 40 45 50 55
 ctt cac cac act ctt atc aag tta gga aag cgt tat ggc cct ttg tac 304
 Leu His His Thr Leu Ile Lys Leu Gly Lys Arg Tyr Gly Pro Leu Tyr
 60 65 70

act ctt tac ttt ggt tcc atg cct acc gtt gtt gca tcc act cct gac	352
Thr Leu Tyr Phe Gly Ser Met Pro Thr Val Val Ala Ser Thr Pro Asp	
75 80 85	
ttg ttt aaa ctt ttc ctt caa acc cat gaa gct act tcc ttt aac aca	400
Leu Phe Lys Leu Phe Leu Gln Thr His Glu Ala Thr Ser Phe Asn Thr	
90 95 100	
aga ttc caa acc tct gct att agt cgt ctt acc tat gac aac tct gtt	448
Arg Phe Gln Thr Ser Ala Ile Ser Arg Leu Thr Tyr Asp Asn Ser Val	
105 110 115	
gct atg gtt cca ttt gca cct tat tgg aag ttt att aga aag ctt atc	496
Ala Met Val Pro Phe Ala Pro Tyr Trp Lys Phe Ile Arg Lys Leu Ile	
120 125 130 135	
atg aac gac ttg ctc aac gcc acc act gtt aac aaa ttg agg cca ttg	544
Met Asn Asp Leu Leu Asn Ala Thr Thr Val Asn Lys Leu Arg Pro Leu	
140 145 150	
agg agc cga gaa atc ctt aag gtt ctt aag gtc atg gct aat agt gct	592
Arg Ser Arg Glu Ile Leu Lys Val Leu Lys Val Met Ala Asn Ser Ala	
155 160 165	
gaa act caa cag cca ctt gat gtc act gag gag ctt ctc aag tgg aca	640
Glu Thr Gln Gln Pro Leu Asp Val Thr Glu Glu Leu Leu Lys Trp Thr	
170 175 180	
aac agc aca atc tct acc atg atg ttg ggt gag gcc gaa gag gtt aga	688
Asn Ser Thr Ile Ser Thr Met Met Leu Gly Glu Ala Glu Glu Val Arg	
185 190 195	
gat att gct cgt gat gtt ctt aag atc ttt gga gaa tat agt gtt aca	736
Asp Ile Ala Arg Asp Val Leu Lys Ile Phe Gly Glu Tyr Ser Val Thr	
200 205 210 215	
aac ttt att tgg cct ttg aac aag ttt aag ttt gga aac tat gat aag	784
Asn Phe Ile Trp Pro Leu Asn Lys Phe Lys Phe Gly Asn Tyr Asp Lys	
220 225 230	
aga act gag gag att ttc aat aag tat gat cct atc att gaa aag gtt	832
Arg Thr Glu Glu Ile Phe Asn Lys Tyr Asp Pro Ile Ile Glu Lys Val	
235 240 245	
atc aag aaa cga caa gag att gtg aac aaa aga aaa aat gga gaa atc	880
Ile Lys Lys Arg Gln Glu Ile Val Asn Lys Arg Lys Asn Gly Glu Ile	
250 255 260	
gta gaa ggc gag cag aat gtt gtt ttt ctt gac act ttg ctt gaa ttt	928
Val Glu Gly Glu Gln Asn Val Val Phe Leu Asp Thr Leu Leu Glu Phe	
265 270 275	
gca caa gat gag acc atg gag atc aaa att aca aag gaa caa atc aag	976
Ala Gln Asp Glu Thr Met Glu Ile Lys Ile Thr Lys Glu Gln Ile Lys	
280 285 290 295	
ggg ctt gtt gtg gat ttt ttc tct gca gga aca gac tcc acc gcc gtg	1024
Gly Leu Val Val Asp Phe Phe Ser Ala Gly Thr Asp Ser Thr Ala Val	
300 305 310	

tct aca gaa tgg act tta tca gag ctc atc aat aat cct aga gtg ttg	1072
Ser Thr Glu Trp Thr Leu Ser Glu Leu Ile Asn Asn Pro Arg Val Leu	
315 320 325	
aag aaa gct cga gag gag att gac tct gtt gtg gga aaa gat aga ctg	1120
Lys Lys Ala Arg Glu Glu Ile Asp Ser Val Val Gly Lys Asp Arg Leu	
330 335 340	
gtt gat gaa tca gat gtt cag aat ctt cct tac att aaa gcc atc gta	1168
Val Asp Glu Ser Asp Val Gln Asn Leu Pro Tyr Ile Lys Ala Ile Val	
345 350 355	
aaa gaa gca ttt cgc ttg cac cca cca cta cct gta gtc aaa aga aaa	1216
Lys Glu Ala Phe Arg Leu His Pro Pro Leu Pro Val Val Lys Arg Lys	
360 365 370 375	
tgt aca caa gaa tgt gag atc gac ggg tat gtg gtt cca gaa gga gca	1264
Cys Thr Gln Glu Cys Glu Ile Asp Gly Tyr Val Val Pro Glu Gly Ala	
380 385 390	
cta ata ctt ttc aat gtc tgg gca gtg gga aga gac cca aaa tat tgg	1312
Leu Ile Leu Phe Asn Val Trp Ala Val Gly Arg Asp Pro Lys Tyr Trp	
395 400 405	
gta aag cca ttg gaa ttt cgt cca gag agg ttc ata gaa aat gtt ggt	1360
Val Lys Pro Leu Glu Phe Arg Pro Glu Arg Phe Ile Glu Asn Val Gly	
410 415 420	
gaa ggt gaa gca gct tca att gat ctt agg ggt caa cat ttc aca ctt	1408
Glu Gly Glu Ala Ala Ser Ile Asp Leu Arg Gly Gln His Phe Thr Leu	
425 430 435	
cta cca ttt ggg tct gga aga agg atg tgt cct gga gtc aat ttg gct	1456
Leu Pro Phe Gly Ser Gly Arg Arg Met Cys Pro Gly Val Asn Leu Ala	
440 445 450 455	
act gca gga atg gcc aca atg att gca tct att atc caa tgc ttc gat	1504
Thr Ala Gly Met Ala Thr Met Ile Ala Ser Ile Ile Gln Cys Phe Asp	
460 465 470	
ctc caa gta cct ggt caa cat gga gaa ata ttg aat ggt gat tat gct	1552
Leu Gln Val Pro Gly Gln His Gly Glu Ile Leu Asn Gly Asp Tyr Ala	
475 480 485	
aag gtt agc atg gaa gag aga cct ggt ctc aca gtt cca agg gca cat	1600
Lys Val Ser Met Glu Glu Arg Pro Gly Leu Thr Val Pro Arg Ala His	
490 495 500	
aat ctc atg tgt gtt cct ctt gca aga gct ggt gtc gca gat aaa ctt	1648
Asn Leu Met Cys Val Pro Leu Ala Arg Ala Gly Val Ala Asp Lys Leu	
505 510 515	
ctt tcc tcc taaaatatct tgagaggatg aatcaccaac atatagcctc	1697
Leu Ser Ser	
520	
tctttggtac tacaaaatta tgatgtaatt ttcttatattt ttctgtcaca aaggaagtgt	1757
tgtaacttgt aattgcatac aaaatctata aattttatca tcctattcat tatt	1811

<210> 5
 <211> 522
 <212> PRT
 <213> Medicago truncatula

 <400> 5
 Met Leu Val Glu Leu Ala Val Thr Leu Leu Leu Ile Ala Leu Phe Leu
 1 5 10 15
 His Leu Arg Pro Thr Pro Thr Ala Lys Ser Lys Ala Leu Arg His Leu
 20 25 30
 Pro Asn Pro Pro Ser Pro Lys Pro Arg Leu Pro Phe Ile Gly His Leu
 35 40 45
 His Leu Leu Asp Asn Pro Leu Leu His His Thr Leu Ile Lys Leu Gly
 50 55 60
 Lys Arg Tyr Gly Pro Leu Tyr Thr Leu Tyr Phe Gly Ser Met Pro Thr
 65 70 75 80
 Val Val Ala Ser Thr Pro Asp Leu Phe Lys Leu Phe Leu Gln Thr His
 85 90 95
 Glu Ala Thr Ser Phe Asn Thr Arg Phe Gln Thr Ser Ala Ile Ser Arg
 100 105 110
 Leu Thr Tyr Asp Asn Ser Val Ala Met Val Pro Phe Ala Pro Tyr Trp
 115 120 125
 Lys Phe Ile Arg Lys Leu Ile Met Asn Asp Leu Leu Asn Ala Thr Thr
 130 135 140
 Val Asn Lys Leu Arg Pro Leu Arg Ser Arg Glu Ile Leu Lys Val Leu
 145 150 155 160
 Lys Val Met Ala Asn Ser Ala Glu Thr Gln Gln Pro Leu Asp Val Thr
 165 170 175
 Glu Glu Leu Leu Lys Trp Thr Asn Ser Thr Ile Ser Thr Met Met Leu
 180 185 190
 Gly Glu Ala Glu Glu Val Arg Asp Ile Ala Arg Asp Val Leu Lys Ile
 195 200 205
 Phe Gly Glu Tyr Ser Val Thr Asn Phe Ile Trp Pro Leu Asn Lys Phe
 210 215 220
 Lys Phe Gly Asn Tyr Asp Lys Arg Thr Glu Glu Ile Phe Asn Lys Tyr
 225 230 235 240
 Asp Pro Ile Ile Glu Lys Val Ile Lys Lys Arg Gln Glu Ile Val Asn
 245 250 255
 Lys Arg Lys Asn Gly Glu Ile Val Glu Gly Glu Gln Asn Val Val Phe
 260 265 270
 Leu Asp Thr Leu Leu Glu Phe Ala Gln Asp Glu Thr Met Glu Ile Lys
 275 280 285

Ile Thr Lys Glu Gln Ile Lys Gly Leu Val Val Asp Phe Phe Ser Ala
 290 295 300
 Gly Thr Asp Ser Thr Ala Val Ser Thr Glu Trp Thr Leu Ser Glu Leu
 305 310 315 320
 Ile Asn Asn Pro Arg Val Leu Lys Lys Ala Arg Glu Glu Ile Asp Ser
 325 330 335
 Val Val Gly Lys Asp Arg Leu Val Asp Glu Ser Asp Val Gln Asn Leu
 340 345 350
 Pro Tyr Ile Lys Ala Ile Val Lys Glu Ala Phe Arg Leu His Pro Pro
 355 360 365
 Leu Pro Val Val Lys Arg Lys Cys Thr Gln Glu Cys Glu Ile Asp Gly
 370 375 380
 Tyr Val Val Pro Glu Gly Ala Leu Ile Leu Phe Asn Val Trp Ala Val
 385 390 395 400
 Gly Arg Asp Pro Lys Tyr Trp Val Lys Pro Leu Glu Phe Arg Pro Glu
 405 410 415
 Arg Phe Ile Glu Asn Val Gly Glu Gly Glu Ala Ala Ser Ile Asp Leu
 420 425 430
 Arg Gly Gln His Phe Thr Leu Leu Pro Phe Gly Ser Gly Arg Arg Met
 435 440 445
 Cys Pro Gly Val Asn Leu Ala Thr Ala Gly Met Ala Thr Met Ile Ala
 450 455 460
 Ser Ile Ile Gln Cys Phe Asp Leu Gln Val Pro Gly Gln His Gly Glu
 465 470 475 480
 Ile Leu Asn Gly Asp Tyr Ala Lys Val Ser Met Glu Glu Arg Pro Gly
 485 490 495
 Leu Thr Val Pro Arg Ala His Asn Leu Met Cys Val Pro Leu Ala Arg
 500 505 510
 Ala Gly Val Ala Asp Lys Leu Leu Ser Ser
 515 520

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/05915

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N15/82 C12N15/11 C12N9/02 C12N5/10
A01H5/00 A01H5/10 A23L1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>SIMINSZKY B. ET AL.: "Cytochrome P450 monooxygenase CYP93C1p from soybean; AC AF022462"</p> <p>EBI DATABASE, 8 January 1998 (1998-01-08), XP002143672 cited in the application the whole document</p> <p>& SIMINSZKY B. ET AL.: "Expression of a soybean cytochrome P450 monooxygenase cDNA in yeast and tobacco enhances the metabolism of phenylurea herbicides" PNAS, U.S.A., vol. 96, February 1999 (1999-02), pages 1750-1755, XP002100400 the whole document</p> <p style="text-align: center;">--- -/--</p>	74-77

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 July 2000

Date of mailing of the international search report

10/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

Int. Jonal Application No

PCT/US 00/05915

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>COLLIVER S P ET AL: "Differential modification of flavonoid and isoflavonoid biosynthesis with an antisense chalcone synthase construct in transgenic Lotus corniculatus."</p> <p>PLANT MOLECULAR BIOLOGY, vol. 35, no. 4, 1997, pages 509-522, XP002143673 ISSN: 0167-4412 the whole document</p> <p>---</p>	1-104
A	<p>PAIVA N L ET AL: "REGULATION OF ISOFLAVONOID METABOLISM IN ALFALFA"</p> <p>PLANT CELL, TISSUE AND ORGAN CULTURE, XX, KLUWER ACADEMIC PUBLISHERS, vol. 38, no. 2/03, 1994, pages 213-220, XP000571330 ISSN: 0167-6857 the whole document</p> <p>---</p>	1-104
A	<p>OTANI K. ET AL.: "Licodione synthase, a cytochrome P-450 monooxygenase catalyzing 2-Hydroxylation of 5-Deoxyflavanone in cultured Glycyrrhiza achinata L. cells"</p> <p>PLANT PHYSIOLOGY, vol. 105, 1994, pages 1427-1432, XP002143674 the whole document</p> <p>---</p>	1-104
A	<p>AKASHI T. ET AL.: "Identification of a cytochrome P450 cDNA encoding (2S)-flavanone 2-hydroxylase of licorice (Glycyrrhiza echinata L.; Fabaceae) which represents licodione synthase and flavone synthase"</p> <p>FEBS LETTERS, vol. 431, 1998, pages 287-290, XP002927621 cited in the application the whole document</p> <p>---</p>	1-104
A	<p>WO 96 15239 A (SAMUEL ROBERTS NOBLE FOUNDATIO) 23 May 1996 (1996-05-23) the whole document</p> <p>---</p>	1-104
A	<p>DIXON RICHARD A ET AL: "The isoflavonoid phytoalexin pathway: From enzymes to genes to transcription factors."</p> <p>PHYSIOLOGIA PLANTARUM, vol. 93, no. 2, 1995, pages 385-392, XP000925614 ISSN: 0031-9317 the whole document</p> <p>---</p>	1-104
	<p>---</p> <p>-/--</p>	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/05915

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DIXON, R. A. ET AL: "Molecular controls for isoflavonoid biosynthesis in relation to plant and human health" 1999, PHYTOCHEMICALS IN HUMAN HEALTH PROTECTION, NUTRITION, AND PLANT DEFENSE, (1999) PP. 133-159. 99 REF. PUBLISHER: KLUWER ACADEMIC/PLENUM PUBLISHERS. NEW YORK ISBN: 0-306-46203-6 XP000925618 see esp. pp.143/44 ---	1-104
P,X	AKASHI TOMOYOSHI ET AL: "Cloning and functional expression of a cytochrome P450 cDNA encoding 2-hydroxyisoflavanone synthase involved in biosynthesis of the isoflavonoid skeleton in licorice." PLANT PHYSIOLOGY (ROCKVILLE), vol. 121, no. 3, November 1999 (1999-11), pages 821-828, XP002143675 ISSN: 0032-0889 see esp. p.1431 r.col. the whole document ---	41
P,X	WO 99 19493 A (UNIV NORTH CAROLINA ;DEWEY RALPH E (US); CORBIN FREDERICK T (US);) 22 April 1999 (1999-04-22) see esp. example 2; SEQ ID NO:9 ---	74-77
P,X	STEELE CHRISTOPHER L ET AL: "Molecular characterization of the enzyme catalyzing the aryl migration reaction of isoflavonoid biosynthesis in soybean." ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, vol. 367, no. 1, 1 July 1999 (1999-07-01), pages 146-150, XP000925681 ISSN: 0003-9861 the whole document ---	74-77
P,X	JUNG WOOSUK ET AL: "Identification and expression of isoflavone synthase, the key enzyme for biosynthesis of isoflavones in legumes." NATURE BIOTECHNOLOGY, vol. 18, no. 2, February 2000 (2000-02), pages 208-212, XP002143676 ISSN: 1087-0156 the whole document -----	1-5, 8-10,15, 16,20, 21, 47-51, 54-56, 74-77, 89, 99-102

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/05915

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9615239 A	23-05-1996	US 5750399 A	12-05-1998
		AU 699496 B	03-12-1998
		AU 3972795 A	06-06-1996
		CA 2204131 A	23-05-1996
		EP 0792356 A	03-09-1997
		JP 10508750 T	02-09-1998
		NZ 296033 A	25-02-1999
WO 9919493 A	22-04-1999	AU 9680698 A	03-05-1999